

2009

Predicting outcome in acute low back pain using different models of patient profiling

Benedict M. Wand

University of Notre Dame Australia, Benedict.Wand@nd.edu.au

James H. McAuley

Louise Marston

Lorraine H. De Souza

Follow this and additional works at: http://researchonline.nd.edu.au/health_article



Part of the [Life Sciences Commons](#), and the [Medicine and Health Sciences Commons](#)

This article was originally published as:

Wand, B. M., McAuley, J. H., Marston, L., & De Souza, L. H. (2009). Predicting outcome in acute low back pain using different models of patient profiling. *SPINE*, 34 (18), 1970–1975.

This article is posted on ResearchOnline@ND at http://researchonline.nd.edu.au/health_article/17. For more information, please contact researchonline@nd.edu.au.



**Predicting outcome in acute low back pain using different models of
patient profiling.**

BM Wand, JH McAuley, L Marston, LH De Souza.

Benedict Martin Wand. PhD

Associate Professor

The University of Notre Dame Australia

School of Health Sciences

19 Mouat St Fremantle

WA 6959, Australia.

James Henry McAuley. PhD

Research Manager

Musculoskeletal Division

The George Institute for International Health

341 George Street

Sydney 2000, Australia

Back Pain Research Group

University of Sydney

East St, Lidcombe

NSW 2141, Australia.

Louise Marston, PhD.

Research Statistician,

Research Department of Primary Care & Population Health,

Division of Population Health,

Faculty of Biomedical Sciences,

University College London Medical School,

London, UK

Lorraine Hillary De Souza. PhD

Chair of Rehabilitation

Department of Health Sciences and Social Care

Brunel University

Kingston Lane, Uxbridge

Middlesex UB8 3PH, UK.

Correspondence to:

Dr Benedict Martin Wand
The University of Notre Dame Australia
School of Health Sciences
19 Mouat St Fremantle
WA 6959, Australia.

bwand@nd.edu.au

Tel +61 8 9433 0203

Fax +61 8 9433 0210

Competing interest: No funds were received in support of this work. No benefits in any form have been or will be received from a commercial party related directly or indirectly to the subjects of this manuscript

Acknowledgements: We would like to thank Christien Bird, Maureen MacDowell, Anne Golden and Mary Sexton for their initial contribution to this study.

Ethical approval: Ethical approval was obtained from the Health Authorities Research Ethics Committee

Abstract

Study Design

Prospective observational study of prognostic indicators, utilising data from a randomised, controlled trial of physiotherapy care of acute low back pain (ALBP) with follow up at 6 weeks, 3 months and 6 months.

Objective

To evaluate which patient profile offers the most useful guide to long-term outcome in ALBP.

Summary of Background Data

The evidence used to inform prognostic decision-making is derived largely from studies where baseline data is used to predict future status. Clinicians often see patients on multiple occasions so may profile patients in a variety of ways. It is worth considering if better prognostic decisions can be made from alternative profiles.

Methods

Clinical, psychological and demographic data were collected from a sample of 54 ALBP patients. Three clinical profiles were developed from information collected at baseline, information collected at 6 weeks, and the change in status between these two time points. A series of regression models were used to determine the independent and relative contributions of these profiles to the prediction of chronic pain and disability.

Results

The baseline profile predicted long-term pain only. The 6-week profile predicted both long-term pain and disability. The change profile only predicted long-term disability ($p < 0.01$). When predicting long-term pain, after the baseline profile had been added to the model, the 6-week profile did not add significantly when forced in at the second step ($p > 0.05$). A similar result was obtained when the order of entry was reversed. When predicting long-term disability, after the 6-week profile was entered at the first step, the change profile was not significant when forced in at the second step. However, when the change profile was entered at the first step and the 6-week clinical profile was forced in at the second step, a significant contribution of the 6-week profile was found.

Conclusions

The profile derived from information collected at 6 weeks provided the best guide to long-term pain and disability. The baseline profile and change in status offered less predictive value.

Key words: acute low back pain; clinical guidelines; prognosis; physiotherapy.

Key points

- International guidelines for ALBP use information about prognosis to shape care pathways for ALBP patients.

- This information is derived largely from studies that have assessed patients at a single (early) time point.
- The clinical situation provides a much richer source of information and potential for varying models of patient profiling.
- The 6-week profile provides the most useful information for predicting long-term outcome.
- On reassessment, the overall status of the patient is a better predictor of outcome than the rate of improvement.

Mini abstract

Guidelines recommend multiple assessments of ALBP patients. We were interested in what information provides the best indicator of chronic status. The 6-week profile was the most useful predictor of long-term status. The baseline profile and change in status offered less predictive value.

Introduction

Low back pain (LBP) is a problem of vast dimensions, it affects up to 80% of the adult population¹ and accounts for considerable healthcare and socioeconomic costs². The scale of the problem has prompted a number of authorities to develop evidence-based guidelines for the management of acute LBP (ALBP)³. These documents provide primary care clinicians with guidance on diagnosis, prognosis and management of the problem based on high quality, clinical research from these three areas.

The information used to provide guidance on prognostic issues has largely been derived from prospective, longitudinal studies where a baseline assessment is made and future clinical status predicted from this single time point^{for e.g.4,5}. The typical clinical experience of managing ALBP provides clinicians with much richer sources of information as patients are generally seen on more than one occasion. Indeed, the algorithms of care that accompany many guidelines promote the idea of serially evaluating the clinical status of patients to determine progression through the algorithm^{for e.g.6}.

Successive patient assessment enables clinicians to formulate impressions of the patient's status based on their initial presentation, subsequent presentations and change status between presentations. It is unclear from the literature which of these three patient profiles is the most useful prognostic model. In order to determine this we decided to conduct a secondary analysis of a randomised, controlled trial of physiotherapy care for ALBP⁷.

Specifically, we were interested in determining if information gathered at baseline or information gathered at an interim follow-up appointment provided the most useful information for predicting long-term pain and disability. We were also interested in determining what information clinicians should attend to at interim appointments. Particularly, whether change in status from baseline or actual status at follow-up was the most useful indicator of long-term clinical outcome. It is hoped that this information will enable primary care clinicians to provide more accurate prognostic information to patients and better inform the decision making process as patients progress through the care pathway.

Materials and Methods

Study participants

This is a secondary analysis of a data set from a randomised, controlled trial of physiotherapy care for acute non-specific low back pain (ANSLBP)⁷. Subjects were 94 ANSLBP patients referred to the Physiotherapy Department of a suburban district hospital in London, England by either their General Practitioner or the Hospital Accident and Emergency Department. To be eligible for inclusion patients had to report non-specific low back pain for less than six-weeks, be aged between 20 and 55 years of age and provide written, informed consent. Those with recurrent pain needed to have been pain free for at least three months prior to the onset of the current episode.

Potential subjects were screened by a physiotherapist for evidence of specific low back pathology (malignancy, fracture, infection, inflammatory disease, etc) or the presence of nerve root pain. Additional exclusion criteria were pregnancy or less than three months post-partum, involvement in litigation related to their back problem, coexisting major medical disease, current involvement in active physical therapy for their problem, or having undergone previous spinal surgery. The study was approved by the Health Authority's Research Ethics Committee.

Procedure

At baseline, subjects completed a set of questions related to their demographic and clinical status. The demographic information collected included, age, gender and work status. The clinical characteristics recorded were duration of the problem and symptom distribution⁸. A screening instrument for psychosocial risk factors, the acute low back pain screening questionnaire (ALBPSQ)⁹, was also administered at baseline.

In addition, patients completed a set of standardised questionnaires that assessed pain, disability, quality of life and psychological functioning. LBP related disability was measured using the Roland and Morris Disability Questionnaire (RMDQ)¹⁰. Pain intensity was calculated by asking subjects to rate their usual pain intensity during the last week on a 0-10 numerical rating scale¹¹. State anxiety was estimated using six items from the Spielberger State-trait Anxiety Inventory (STAI)¹². The presence of depressive symptoms was

determined using the Modified Zung Self Rated Depression Score (Zung)¹³, and distress was estimated using the Modified Somatic Perception Questionnaire (MSPQ)¹⁴. Quality of life was measured using the EuroQol health transition score (EQ5D)¹⁵, Physical well-being was calculated from the Short Form-36 physical component score (PCS)¹⁶ and mental well-being from the Short Form-36 mental component score (MCS)¹⁶. All patients completed these questionnaires at baseline and were resent the assessments at six-weeks, three-months and six-months.

Predictor variables

All variables measured at baseline (shown in table 1.) were used as predictor variables. The six-week scores for Pain, RMDQ, STAIS, Zung, MSPQ, EQ5D, PCS and MCS were also used as predictor variables. Change scores were calculated by subtracting the six-week scores from the baseline scores for those variables that were measured at these two time points, giving each patient a value that represented the relative amount of change, these change scores were also included as predictor variables. Predictor variables measured at baseline formed the *acute clinical profile*, those measured at six-weeks formed the *sub-acute clinical profile* and the change scores were used to determine the *change clinical profile*.

Outcome

The outcomes of interest were long-term back pain related disability and long-term pain intensity. These were derived from the mean scores of the three and

six month assessments of the RMDQ and the usual pain intensity numerical rating scale respectively.

Data analysis

Predictor variables that demonstrated significant bivariate correlations (Pearson's r) with long-term disability and long-term pain were identified and classified into their respective acute, sub-acute and change clinical profiles. The significance level was set at $p < 0.01$ to account for multiple comparisons.

A series of multiple regression models were fit to determine the independent contribution of the acute, sub-acute and change profiles to the prediction of long-term disability and long-term pain. The relative contribution of the clinical profiles to the two outcomes was determined by a series of hierarchical regressions models where the order of entry of the profiles was rotated. All analyses were undertaken using SPSS for windows version 15.

Results

Full data was available for 54 patients. The baseline demographic and clinical characteristics of responders and non-responders are presented in table 1. There were no significant differences in baseline values between those patients who provided complete data at all time points and those who did not ($p > 0.05$).

Correlation summary

The variables that had significant Pearson's correlations ($p < 0.01$) with either long-term pain or long-term disability are presented in table 2, classified into their respective clinical profiles.

Regression models

The regression models showing the relationships between the clinical profiles and long-term pain and disability are shown in table 3. This demonstrates that the sub acute ($R^2 = 0.607$) and change ($R^2 = 0.131$) profiles were associated with long-term disability and the acute ($R^2 = 0.159$) and sub-acute profiles ($R^2 = 0.257$) were associated with long-term pain.

The results of the hierarchical regression model with long-term pain as the dependent variable showed that when the acute clinical profile was entered at the first step, the sub-acute profile was not significant when forced into the model at the second step ($p > 0.05$). A similar result was obtained when the order was reversed.

The result of the hierarchical regression model with long-term disability as the dependent variable showed that when the sub-acute profile was entered at the first step, the change profile was not significant when forced into the model at the second step. However, when the change profile was entered at the first step and the sub-acute profile was forced in at the second step, a significant contribution of the sub-acute profile was demonstrated (R^2 change = 0.486; F change = 15.203; $df = 4, 48$; $p < 0.001$).

These results indicate that the sub-acute profile provides the most valuable information for predicting long-term disability. Some useful information on long-term pain may be obtained from the acute and sub-acute profiles, however it appears the sub acute profile has stronger predictive value

Discussion

Summary of main findings

Clinicians have been encouraged to consider the acute patient profile in treatment planning and prognostic decision-making. Despite a comprehensive baseline profile of patients with ALBP, we found very little of interest in predicted chronic status. No baseline variable was predictive of long-term disability and only the ALBPSQ score was predictive of long-term pain. Notably, no uni-dimensional estimate of patients' acute psychological function appeared to impact on long-term outcome.

We were interested in whether other information may be useful to clinicians and found that the sub-acute clinical profile and the short-term rate of change provided some information on who may develop chronic symptoms. The sub-acute profile appears to be more meaningful. Measures of sub-acute pain intensity, disability (RMDQ), physical well being (PCS), mood (Zung) and general health (EQ5D) were predictive of long-term disability and together explained over 60% of the variance. Only pain intensity, disability (RMDQ) physical well being (PCS), and general health (EQ5D) were useful predictors of chronic pain, and the combined explanatory power was significantly less (26%).

The change in disability (RMDQ) was significantly associated with chronic disability and explained about 13% of the variance. No change variable was significantly related to long-term pain. Change in clinical status is only marginally useful in predicting chronic disability, and of no value in predicting chronic pain. This finding was contrary to our expectations. We had anticipated that patients who demonstrated large changes in their clinical profile would have favourable outcomes. These data suggest that on reassessment the overall status of the patient is a better predictor of outcome than the rate of improvement.

We conducted a series of multivariate analyses to try to discern the *relative* importance of the different clinical profiles. These analyses demonstrated that the sub-acute profile contains the most unique information for predicting long-term disability, providing considerable information above that which is derived from change status. When predicting long-term pain, the acute and sub-acute profiles provide equally important information.

These results highlight the complex relationship between pain and disability. The clinical features that predict chronic pain and disability vary and the explanatory power is very different. When seeking information on prognosis it is important that clinicians are clear on what outcomes are of interest to them and their patients and at what stage the patient is when making this decision.

Strengths and limitations

There are several strengths to this research. We used a comprehensive set of assessments which sampled pain, disability, psychological function and health related quality of life, measured on the same cohort of patients, longitudinally. Furthermore, data were collected in the clinical environment, which reflects the reality of day-to-day clinical practice.

The main limitations include the small number of subjects and the proportion of patients who did not provide full follow-up data and were therefore excluded from the analyses. The sample size is small for the number of statistical tests undertaken, however, we have attempted to control for this by adopting a more stringent significance level. Furthermore, patients who did not provide full follow up data and were excluded from the analysis did not have significantly different initial presentations from those who provided complete data at all time points (table 1.). While this analysis indicates that the data may be missing at random, care must always be taken when interpreting results with this level of loss to follow up.

Additionally, this study was performed within the framework of a randomised controlled trial potentially lowering the external validity for answering prognostic questions. All outcomes used were self-reported measures and may be biased by some shared method variance¹⁷. Finally, as with all prognostic research, our models may be limited by not having measured adequate prognostic factors. Our findings should be interpreted with some caution and our prognostic models now require testing in large-scale prospective clinical studies.

Comparison with existing literature

Our results support earlier work that suggests the ALBPSQ has some value in predicting chronic status in ALBP patients^{9,18,19,20}. It appears that information about long-term pain levels can be obtained from multidimensional evaluation of psychosocial status at baseline. Other researchers have noted that some uni-dimensional measures of psychosocial status are also predictive of outcome. Job dissatisfaction²¹, previous sick leave for LBP⁵, somatic distress^{22,23}, depression^{24,25,26 27,28,29}, fear of movement¹⁷ and passive coping^{23,30} have all been shown to predict long-term status when measured at baseline. We assessed patients' anxiety, somatic distress, depression and mental well-being at baseline and found little of importance in determining long-term pain or disability with these measures.

Some of this discrepancy may lie in the timing of clinical evaluation. The present study only sourced patients whose current episode was less than six weeks, and the average time since onset of the baseline assessment was less than three weeks. Studies that have found depression a useful predictor have used a less strict inception cohort^{24,25,26,29} or collected data sometime after the initial consultation²⁷. In support of this view, we found that depression measured at six weeks was significantly correlated with chronic disability. It may be that high levels of depressive symptoms in the very acute phase are less important, maintenance of depression into the sub-acute phase or development at the sub-acute phase might be the primary problem.

A common finding in prognostic studies on ALBP is the relationship between high pain intensity at baseline and future status³¹. Our analyses found no relationship between baseline pain and either chronic pain or chronic disability. We noted a similar trend to that seen for depressive symptoms. Pain levels measured acutely were not related to chronic outcome, though sub-acute measures of pain were correlated with long-term pain and disability. The explanation may be a reflection of the small sample size in the current study, or lie in mixed populations^{23,22} and different inception cohorts^{24,30,32} used by other investigators. In support of this, the systematic review by Pengel et al.³³ reviewed only papers with an inception cohort of less than three weeks and did not find pain intensity a useful predictor of outcome.

Other groups have also noted improvement in prognostic accuracy with repeated assessment. Enthoven et al.³⁴ performed a series of physical tests on a group of patients with LBP of varying duration at initial presentation and again four weeks later. They found none of the physical measures at baseline to be associated with long-term disability, yet three of the four measures taken at week four were related to disability at 12 months. Klenerman et al.³⁵ assessed patients at one week and two months. The two month data explained considerably more of the variance in 12 month outcome than data collected at week one. Likewise Carey et al.³⁶ found week four assessment of functional status a far stronger predictor of chronic outcome than baseline assessment. Heneweer and colleagues³⁷ dichotomised patients into recovered and not recovered at 12 weeks. They noted no difference in pain and disability between these two groups at the two-week assessment. However, they were

clearly delineable at the four and eight week assessments. These results should perhaps not be surprising, as the more delayed assessment profiles the patient at a time closer to the final evaluation.

Dunn and Croft³⁸ undertook a detailed analysis of this phenomenon on a group of predominantly chronic LBP (CLBP) patients. Their results clearly demonstrate that repeat assessment of patients enables a more accurate prediction of prognosis. The analyses used included classifying patients based on the stability of clinical characteristics between the two time points. They showed that people who have persistence of prognostic indicators had the greatest risk of poor outcome. Finally, Sieben et al.³⁹ saw a slightly different pattern in a group of ALBP patients who were monitored daily for two weeks. This study found rising levels of pain-related fear, rather than stable levels, were a stronger predictor of outcome. We found the change in status to be less informative than actual sub-acute status and the hierarchical regression analysis demonstrated that the change profile did not significantly improve the explanatory power of the sub-acute profile. Further work is needed to ascertain the most meaningful information that can be extracted from serial evaluation and whether this differs between acute and chronic patients.

Conclusion

The usefulness of clinical information in making decisions about prognosis in ALBP patients is influenced by the time at which it is collected and the outcome of interest. The useful predictors of long-term pain and disability are different and the variance that can be explained is quite disparate. When

serially assessing ALBP patients, clinicians may obtain more accurate information about long-term outcome from follow-up assessments. Furthermore, the actual status at follow-up appears to be a much more useful guide to long-term outcome than the amount of change in status from baseline.

References

1. Walker BF, Muller R, Grant WD. Low back pain in Australian adults: prevalence and associated disability. *Journal Of Manipulative And Physiological Therapeutics* 2004;27:238-244.
2. Maniadakis N, Gray A. The economic burden of back pain in the UK. *Pain* 2000;84:95-103.
3. Koes BW, van Tulder MW, Ostelo R, et al. Clinical guidelines for the management of low back pain in primary care: an international comparison. *Spine* 2001;26:2504.
4. Coste J, Delecoeuillerie G, Cohen de Lara A, et al. Clinical course and prognostic factors in acute low back pain: an inception cohort study in primary care practice. *BMJ* 1994;308:577-580.
5. Schiottz-Christensen B, Nielsen GL, Hansen VK, et al. Long-term prognosis of acute low back pain in patients seen in general practice: a 1-year prospective follow-up study. *Family Practice* 1999;16:223-232.
6. Chou R, Qaseem A, Snow V, et al. Diagnosis and treatment of low back pain: a joint clinical practice guideline from the American College of Physicians and the American Pain Society. *Annals Of Internal Medicine* 2007;147:478-491.
7. Wand BM, Bird C, McAuley JH, et al. Early intervention for the management of acute low back pain: a single-blind randomized controlled trial of biopsychosocial education, manual therapy, and exercise. *Spine* 2004;29:2350-2356.

8. Spitzer W, LeBlanc F, Dupuis M (eds). Scientific approach to the assessment and management of activity-related spinal disorders: A monogram for clinicians. Report of the Quebec Task Force on Spinal Disorders. *Spine* 1987;12:S1-S60.
9. Linton SJ, Hallden K. Can we screen for problematic back pain? A screening questionnaire for predicting outcome in acute and subacute back pain. *The Clinical Journal Of Pain* 1998;14:209-215.
10. Roland M, Morris R. A study of the natural history of back pain. Part I: development of a reliable and sensitive measure of disability in low-back pain. *Spine* 1983;8:141-144.
11. Bolton JE. Accuracy of recall of usual pain intensity in back pain patients. *Pain* 1999;83:533-539.
12. Spielberger CD, Gorsuch RL, Lushene R. *The State-trait Anxiety Inventory Manual*. Palo Alto, California: Consulting Psychologists Press; 1970.
13. Main CJ, Waddell G. The detection of psychological abnormality in chronic low back pain using four simple scales. *Current Concepts in Pain* 1984;2:10-15.
14. Main CJ. The Modified Somatic Perception Questionnaire (MSPQ). *Journal Of Psychosomatic Research* 1983;27:503-514.
15. The EuroQol Group. EuroQol - a new facility for the measurement of health-related quality of life. *Health Policy* 1990;16:199-208.
16. Ware JE, Snow KK, Gandek B. *SF-36 Health Survey Manual and Interpretation Guide*. Boston: New England Medical Center, The Health Institute; 1993.

17. Swinkels-Meewisse IEJ, Roelofs J, Schouten EGW, et al. Fear of movement/(re)injury predicting chronic disabling low back pain: a prospective inception cohort study. *Spine* 2006;31:658-664.
18. Linton SJ, Boersma K. Early identification of patients at risk of developing a persistent back problem: the predictive validity of the Orebro Musculoskeletal Pain Questionnaire. *The Clinical Journal Of Pain*. 2003;19:80-86.
19. Grotle M, Brox JI, Glomsrod B, et al. Prognostic factors in first-time care seekers due to acute low back pain. *European Journal Of Pain* 2007;11:290-298.
20. Hockings RL, McAuley JH, Maher CG. A systematic review of the predictive ability of the Orebro Musculoskeletal Pain Questionnaire. *Spine* 2008;33:E494-E500.
21. Coste J, Lefrancois G, Guillemin F, et al. Prognosis and quality of life in patients with acute low back pain: insights from a comprehensive inception cohort study. *Arthritis And Rheumatism* 2004;51:168-176.
22. Grotle M, Brox JI, Veierod MB, et al. Clinical course and prognostic factors in acute low back pain: patients consulting primary care for the first time. *Spine* 2005;30:976-982.
23. Burton AK, Tillotson KM, Main CJ, et al. Psychosocial predictors of outcome in acute and subchronic low back trouble. *Spine* 1995;20:722-728.
24. Cherkin DC, Deyo RA, Street JH, et al. Predicting poor outcomes for back pain seen in primary care using patients' own criteria. *Spine* 1996;21:2900-2907.

25. Epping-Jordan JE, Wahlgren DR, Williams RA, et al. Transition to chronic pain in men with low back pain: predictive relationships among pain intensity, disability, and depressive symptoms. *Health Psychology* 1998;17:421-427.
26. Thomas E, Silman AJ, Croft PR, et al. Predicting who develops chronic low back pain in primary care: a prospective study. *BMJ* 1999;318:1662-1667.
27. Dionne CE. Psychological distress confirmed as predictor of long-term back-related functional limitations in primary care settings. *Journal Of Clinical Epidemiology* 2005;58:714-718.
28. Sieben JM, Vlaeyen JWS, Portegijs PJM, et al. A longitudinal study on the predictive validity of the fear-avoidance model in low back pain. *Pain*. 2005;117:162-170.
29. Young Casey C, Greenberg MA, Nicassio PM, et al. Transition from acute to chronic pain and disability: a model including cognitive, affective, and trauma factors. *Pain*. 2008;134:69-79.
30. Jones GT, Johnson RE, Wiles NJ, et al. Predicting persistent disabling low back pain in general practice: a prospective cohort study. *The British Journal Of General Practice* 2006;56:334-341.
31. Kent PM, Keating JL. Can we predict poor recovery from recent-onset nonspecific low back pain? A systematic review. *Manual Therapy*. 2008;13:12-28.
32. Bekkering GE, Hendriks HJM, van Tulder MW, et al. Prognostic factors for low back pain in patients referred for physiotherapy:

- comparing outcomes and varying modeling techniques. *Spine*. 2005;30:1881-1886.
33. Pengel LHM, Herbert RD, Maher CG, et al. Acute low back pain: systematic review of its prognosis. *BMJ* 2003;327:323-323.
 34. Enthoven P, Skargren E, Kjellman G, et al. Course of back pain in primary care: a prospective study of physical measures. *Journal Of Rehabilitation Medicine* 2003;35:168-173.
 35. Klenerman L, Slade PD, Stanley IM, et al. The prediction of chronicity in patients with an acute attack of low back pain in a general practice setting. *Spine* 1995;20:478-484.
 36. Carey TS, Garrett JM, Jackman AM. Beyond the good prognosis. Examination of an inception cohort of patients with chronic low back pain. *Spine* 2000;25:115-120.
 37. Heneweer H, Aufdemkampe G, van Tulder MW, et al. Psychosocial variables in patients with (sub)acute low back pain: an inception cohort in primary care physical therapy in The Netherlands. *Spine* 2007;32:586-592.
 38. Dunn KM, Croft PR. Repeat assessment improves the prediction of prognosis in patients with low back pain in primary care. *Pain* 2006;126:10-15.
 39. Sieben JM, Vlaeyen JWS, Tuerlinckx S, et al. Pain-related fear in acute low back pain: the first two weeks of a new episode. *European Journal Of Pain* 2002;6:229-237.

Table 1: Comparison of baseline status between those included (responders) and excluded (non responders) from analysis

<i>Variables</i>	<i>Responders (n=54)</i>		<i>Non Responders (n=40)</i>		p-value
	N or mean	% or (SD)	N or mean	% or (SD)	
Age (range)	35 (21, 55)	(9)	34 (21, 52)	(8)	0.616
Male	26	48	21	53	0.677
BMI	25	(4)	26	(4)	0.565
Symptom distribution					
No symptoms	0	0	1	3	0.724
LBP without radiation	30	56	22	55	
Proximal radiation	12	22	7	18	
Distal radiation	12	22	10	25	
Uses analgesics	31	57	21	53	0.636
Duration (weeks)	2.9	(1.4)	3.0	(1.5)	0.766
Work status					
Off work	22	41	16	40	0.480
Working	28	52	18	45	
Not employed	4	7	6	15	
ALBPSQ	89	(27)	95	(31)	0.307
PCS	36	(7)	38	(7)	0.305
MCS	48	(8)	46	(9)	0.213
EQ5D	0.60	(0.25)	0.57	(0.28)	0.590
Pain	5.2	(2.3)	5.8	(2.2)	0.195
RMDQ	11	(6)	12	(6)	0.565
Zung	21	(10)	23	(12)	0.286
MSPQ	7.3	(5.3)	7.5	(5.0)	0.843
STAIS	13	(4)	13	(4)	0.973

BMI indicates body mass index. ALBPSQ, acute low back pain screening questionnaire (possible range 2-210). PCS, SF36 physical component score (possible range 0-100). MCS, SF36 mental component score (possible range 0-100). EQ5D, EuroQol health transition score (possible range -0.59-1). Pain, numerical rating scale for usual pain intensity (possible range 0-10). RMDQ, Roland and Morris disability questionnaire (possible range 0-24). Zung, Modified Zung self reported depression scale (possible range 0-69). MSPQ, Modified somatic perceptions questionnaire (possible range 0-39). STAIS, Spielberger state-trait anxiety inventory score (possible range 6-24).

Table 2. Correlation coefficients for predictor variables that were significantly related to long-term outcome ($p < 0.01$), classified into their respective clinical profiles.

	<i>r long-term disability</i>	<i>r long term pain</i>
acute profile		
ALBPSQ	0.34*	0.40
subacute profile		
Pain	0.50	0.40
RMDQ	0.73	0.48
PCS	-0.46	-0.36
EQ5D	-0.70	-0.42
Zung	0.45	0.11*
change profile		
RMDQ	0.36	0.12*

*correlations were not significant $p < 0.01$.

Table 3: Results of the multiple regression models of the three clinical profiles on the dependent variables of long-term pain and disability

<i>clinical profiles</i>	<i>dependent variable</i>	<i>R</i>	<i>R²</i>	<i>Ad R²</i>	<i>F</i>	<i>df</i>	<i>Sig F change</i>
acute	<i>Long-term RMDQ</i>	*					
	<i>Long-term Pain</i>	0.398	0.159	0.143	9.809	1, 52	0.003
subacute	<i>Long-term RMDQ</i>	0.779	0.607	0.566	14.809	5, 48	<0.001
	<i>Long-term Pain</i>	0.507	0.257	0.196	4.237	4, 49	0.005
change	<i>Long-term RMDQ</i>	0.362	0.131	0.114	7.843	1, 52	0.007
	<i>Long-term Pain</i>	*					

*No correlations between the outcome and any variables from those clinical profiles had $p < 0.01$, so no regressions were carried out.