2008

Progress in the Fremantle Primary Prevention Study- A Pilot Study of Risk Factor Modification for Cardiovascular Disease

Diane Arnold-Reed
University of Notre Dame Australia, darnold-reed@nd.edu.au

Tom Brett
University of Notre Dame Australia, tom.brett@nd.edu.au

W. Walker

Frances Cadden
University of Notre Dame Australia, Frances.Cadden@nd.edu.au

J. Young

See next page for additional authors

This conference paper was originally published as:
Arnold-Reed, D., Brett, T., Walker, W., Cadden, F., Young, J., Mora, N., Manea-Walley, W., Hince, D., & Caneppele, M. (2008). Progress in the Fremantle Primary Prevention Study- A Pilot Study of Risk Factor Modification for Cardiovascular Disease. *2008 GP & PHC Research Conference: Health for All?*.

Authors
Diane Arnold-Reed, Tom Brett, W. Walker, Frances Cadden, J. Young, Noelene Mora, Wendy Manea-Walley, Dana Hince, and M. Caneppele

Fremantle Primary Prevention Study

A pilot study of risk factor modification for cardiovascular disease

Arnold-Reed D[1], Brett T[1], Hince D[1], Bulsara M[1,2], Sotzik F[1], Caneppele M[1] & O'Driscoll G[1].

[1]General Practice and Primary Health Care Research, School of Medicine, University of Notre Dame Australia, Fremantle
[2]School of Population Health, Faculty of Medicine, Dentistry and Health Sciences, University of Western Australia

Coolibah Medical Centre Greenwood
GP: Dr Frances Cadden
Practice Nurse: Wendy Manea-Walley

Mosman Park Medical Group
GP: A/Prof Tom Brett
Practice Nurse: Julie Young

Murray Medical Centre Mandurah
GP: Dr William Walker
Practice Nurse: Noelene Mora

Background

Risk factors for cardiovascular disease (obesity, diabetes, hypertension, smoking, inactivity, hyperlipidemia) are modifiable, but often tend to be taken seriously only AFTER a significant event occurs (heart attack, stroke).

General practice offers the opportunity to detect and attempt to modify these risk factors and help reduce subsequent morbidity and mortality.

Objectives

Document the prevalence of modifiable cardiovascular risk factors in 40 – 80 year old men and women.

Monitor the changes in cardiovascular risk following delivery of intensive and opportunistic intervention strategies at the GP level.

Methods



Results/Discussion

To date, 559 participants have completed the study (intensive - n=155, 75 ♀; opportunistic - n=404, 161 ♀).

The figures below represent % of participants outside target for key outcome variables for those that have completed the 12 month study.





FIGURE NOTE: *p<0.05, **p<0.01, ***p<0.001; McNemara χ^2 using the binomial distribution

Implications for policy & practice

While a significant proportion of the community are already known to have cardiovascular risk factors, GPs and their practice staff are ideally positioned to assess the extent of this potential morbidity and institute treatment modalities and programmes to reverse and modify them.

Acknowledgement: This study was funded by Pfizer AUS-Non Drug CVS-06-001