

2009

Fitness, motor competence and body composition are weakly associated with adolescent back pain

Mark Perry

Curtin University of Technology, Mark.Perry@curtin.edu.au

Leon Straker

Curtin University of Technology, L.Straker@curtin.edu.au

Peter O'Sullivan

Curtin University of Technology, P.OSullivan@curtin.edu.au

Anne Smith

Curtin University of Technology, Anne.Smith@exchange.curtin.edu.au

Beth P. Hands

University of Notre Dame Australia, bhands@nd.edu.au

Follow this and additional works at: https://researchonline.nd.edu.au/health_article



Part of the [Life Sciences Commons](#), and the [Medicine and Health Sciences Commons](#)

This article was originally published as:

Perry, M., Straker, L., O'Sullivan, P., Smith, A., & Hands, B. P. (2009). Fitness, motor competence and body composition are weakly associated with adolescent back pain. *Journal of Orthopedic and Sports Physical Therapy*, 39 (6), 439-449.

<http://doi.org/10.2519/jospt.2009.3011>

This article is posted on ResearchOnline@ND at https://researchonline.nd.edu.au/health_article/27. For more information, please contact researchonline@nd.edu.au.



1 **Title:** Fitness, motor competence and body composition are weakly associated with
2 adolescent back pain

3
4

5 **ABSTRACT**

6
7

Study Design

8 Cross-sectional survey

9 **Objectives**

10 The objective of this study was to assess the associations between adolescent back pain and
11 fitness, motor competence and body composition.

12 **Background**

13 Although deficits in physical fitness and motor control have been shown to relate to adult back
14 pain, the evidence in adolescents is less clear.

15 **Methods**

16 In this cross-sectional study, 1608 'Raine' cohort adolescents (mean age 14 years) answered
17 questions on lifetime, month and chronic prevalence of back pain, and participated in a range
18 of physical tests assessing aerobic capacity, muscle performance, flexibility, motor
19 competence and body composition. A history of any diagnosed back pain in the adolescent
20 was obtained from the primary carer.

21 **Results**

22 After multivariate logistic regression analysis, increased likelihood of back pain in boys was
23 associated with greater aerobic capacity, greater waist girth and both reduced and greater
24 flexibility. Back pain in girls was associated with greater abdominal endurance, reduced
25 kinaesthetic integration, and both reduced and greater back endurance. Lower likelihood of

1 back pain was associated with greater bimanual dexterity in boys and greater leg power in
2 girls.

3

4 **Conclusion**

5 Physical characteristics are commonly cited as important risk factors in back pain
6 development. Although some factors were associated with adolescent back pain, and these
7 differed between boys and girls, they made only a small contribution to logistic regression
8 models for back pain. The results suggest future work should explore the interaction of
9 multiple domains of risk factors (physical, lifestyle and psychosocial) and subgroups of
10 adolescent back pain, for whom different risk factors may be important.

11

12 **Key words:** Spinal pain; physical performance; motor control; Raine

13

14

15

1 INTRODUCTION

2 Over half of adolescents may have had back pain at some point in their lives¹⁴, one third of
3 these having sought professional help¹¹ and 20% having experienced a reduced quality of
4 life¹¹. The identification of risk factors to facilitate effective prevention and better
5 management of adolescent back pain is therefore important, particularly as adolescent back
6 pain has also been associated with future back pain¹¹. Previous research into adolescent back
7 pain has established that certain lifestyle and psychosocial factors such as computer use³⁷ or
8 poor mental health⁴⁰ are important, but the contribution of physical risk factors in adolescence
9 is still unclear.

10
11 In adults, obesity⁴, deficits in aerobic fitness²⁷, poorer muscle performance^{4,5,27} and reduced
12 motor control⁸ are established physical risk factors, and interventions for adult back pain
13 aimed at countering spinal deconditioning reflect this¹². Such factors may render the spine
14 vulnerable to tissue strain and pain²⁸, and reinforce the pain avoidance / depression cycle³⁹.
15 However, the association between such physical factors and back pain may be different in
16 adolescence, as a result of factors such as the growth spurt, which is known to lead to
17 musculoskeletal changes^{18,19}.

18
19 Current evidence of a relationship between adolescent back pain and physical risk factors is
20 conflicting or limited. Adolescent back pain has been associated with increased adiposity in
21 some studies (e.g.^{19,35}) but not others (e.g.^{16,31}). No studies have examined the relationship
22 between objectively measured aerobic capacity and adolescent back pain, although a recent
23 study of adolescents found aerobic capacity to be unrelated to undifferentiated neck/back
24 pain³. Reduced trunk muscle endurance has been shown to be a risk factor (e.g.^{16,32}), but not
25 by all studies³¹, and no studies have investigated the influence of limb muscle performance.

1 Reduced lumbar or hamstring flexibility has been shown to relate to back pain in some studies
2 (e.g.^{16,19,32,35}), but not by all^{16,24,31}. Finally, the association between motor competence and
3 adolescent back pain is yet to be reported, although pre-pubertal children with lower motor
4 competence report more back pain⁷.

5
6 Differences in study design or definitions of back pain across different studies may partially
7 explain discrepancies between studies. For example, back pain was variously defined as the
8 history of at least one episode¹⁶, pain lasting more than a day²⁴, pain interfering with function
9 for at least one week¹⁹, or pain that did not include menstrual or traumatic pain³⁵. Failure to
10 show effects in some studies may also be due to limitations such as insufficient sample size,
11 with most having samples of <100 (e.g.^{16,19,31,32,35}). One drawback of all the previous literature
12 into physical factors and back pain has been to consider only relationships modelled by a
13 straight line (rectilinear), which may fail to identify those more appropriately modelled by a
14 regular curve (curvilinear), despite reports of curvilinear relationships between spinal pain and
15 activity⁴⁰ and computer use³⁷. Another potential limitation has been inadequate multivariate
16 analysis. Though most studies have looked at several variables, these have mostly, within a
17 given study, focussed on only one or two domains of physical fitness, such as trunk muscle
18 performance³¹ or flexibility^{19,31}. It is therefore unclear whether variables associated with pain
19 are merely correlates of other (possibly more clinically relevant) aspects that have not been
20 considered.

21
22 Therefore the aim of this study was to investigate, within a large cohort, the relationships
23 between a broad range of physical risk factors (body composition, aerobic fitness, muscle
24 performance, flexibility, motor competence), allowing for curvilinear relationships and
25 different risk factor relationships for adolescent boys and girls.

1 **METHODS**

2 **Design**

3 This was a cross-sectional epidemiological study. The study was approved by the Human
4 Research Ethics Committees of Curtin University of Technology and Princess Margaret
5 Hospital. Adolescents provided written informed assent and their parent/guardian provided
6 written informed consent prior to participation. The rights of all participants were protected.

7

8 **Participants**

9 Data from 1608 adolescents (783 girls, 825 boys) of mean (SD) age 14.06 (0.20) yrs were
10 collected as part of their participation in the Western Australian Pregnancy Cohort “Raine”
11 Study (www.rainestudy.org.au). This long term project started with a cohort of women
12 attending antenatal clinics at King Edward Memorial Hospital for Women, Perth, Australia
13 between 1989 and 1991. The children have been followed at birth, 1, 2, 3, 5, 8, 10, and now 14
14 years of age. Inclusion criteria for the women included gestational age between 16 and 20
15 weeks, sufficient proficiency in English to understand the implications of participation, and an
16 intention to remain in Western Australia so that follow-up would be possible. There were 2337
17 adolescents eligible for the 14 year follow-up, of which 1704 (73%) consented to some aspect
18 of the follow-up and 1608 (69%) completed the data collection requirements for the analysis
19 reported in this paper. There were no exclusion criteria for this follow up cohort.

20 **Outcome measures**

21 Participants completed a questionnaire on a laptop at an assessment centre with the help of a
22 research assistant. The questionnaire contained 130 questions concerning a broad range of
23 physical, medical, nutritional, psychosocial and developmental issues. The back pain questions
24 were: Have you ever had back pain? (“yes” or “no”), Has your back been painful in the last
25 month? (“yes” or “no”), and Did your back pain last for more than 3 months? (“yes” or “no”).

1 Prior questions (not relevant to this report) on neck pain and limb pain alerted participants that
2 “back pain” did not include neck or limb pain. The full questionnaire took about 1 hour to
3 complete, and the back pain questions occurred in the first half. Similar versions of these
4 questions have been validated¹⁴.

5
6 Information on diagnosed back pain was obtained from the primary carer, who was asked,
7 “Does your child have now, or has your child had in the past, any of the following health
8 professional diagnosed medical conditions or health problems?”. The primary carer had to
9 indicate which medical diagnoses their child had experienced from a short list of general
10 medical problems, which included “back pain”. This question was part of a questionnaire
11 given to the primary carer, covering many other factors not relevant to this report.

12
13 A physical assessment of the child was carried out after the questionnaire. All tests were
14 carried out by trained and experienced graduate research assistants with a nursing or human
15 movement background. With shoes removed, height (m) was measured with a stadiometer,
16 body mass (kg) with digital scales, waist girth (cm) was measured at the umbilical level with a
17 cloth tape, and arm girth (cm) was measured at the mid-humeral point with a cloth tape. A
18 series of physical performance tests were then conducted, all of which have been previously
19 validated in very similar forms^{6,20,22,25,30,34,38}. Reliability of comparable forms of these tests is
20 also good^{15,20,21,22,25,30} though there are no reports on the reliability of the basketball throw.
21 Most of these validity and reliability studies were conducted on pre-adults^{15,22,30,34,38}. These
22 tests are described as follows.

23
24 Maximal aerobic capacity was estimated using heart rate recordings during sub-maximal cycle
25 ergometry using the Physical Work Capacity 170 protocol¹⁰. Trunk endurance was assessed

1 by the sustained back extension test⁵ and the number of abdominal curls performed in 3
2 minutes¹. Limb muscle performance was evaluated by standing long jump²², seated basketball
3 throw¹ and grip strength²². Hamstring flexibility was tested using a unilateral sit and reach
4 test¹. Finally, motor competence was assessed using the McCarron Assessment of
5 Neuromuscular Development (MAND)²². This assessment measures sensorimotor
6 neuromuscular development normalised to age, and an overall score between 0 (poorest) and
7 100 is obtained - the Neuromuscular Developmental Index (NDI)²². The NDI score is based
8 on performance in ten sensorimotor tests, and eight of these make up the 4 sub-indices of
9 Bimanual Dexterity, Muscle Power, Kinaesthetic Integration and Persistent Control (Table 1).
10 Bimanual Dexterity measures co-ordination across the upper limbs, Muscle Power measures
11 upper limb strength and lower limb power, Kinaesthetic Integration measures gross balance,
12 and Persistent Control measures the ability to produce smooth controlled movements²².

13

14 Insert Table 1 about here

15

16 **Data analysis**

17 All statistical analysis was carried out using SPSS version 15 (SPSS Inc., Chicago, USA).
18 Gender differences were analysed using independent t tests for each of the continuous
19 variables, and Chi squared tests for the categorical variables. To facilitate the interpretation of
20 non-linear relationships, continuous variables were categorised into the bottom 25%, inter-
21 quartile range and top 25%, and the proportions of subjects with back pain in these categories
22 were compared.

23

24 Univariate logistic regression models predicting lifetime, last month, chronic (lasting more
25 than 3 months) and diagnosed back pain from each physical performance characteristic were

1 calculated, with statistical significance set at $p < 0.05$, and the interquartile range (IQR) of
2 each continuous variable (the range between the 25th and 75th percentiles) was defined as the
3 reference category. Corrections for multiple univariate tests were not performed as the
4 multivariate results were the end point of the study. Male and female data were analysed
5 independently as a previous study has reported significantly different physical performance
6 between genders¹³.

7
8 Backwards stepwise likelihood ratio multivariate logistic regression models were used to
9 evaluate the combined associations of performance factors for boys and for girls separately,
10 with the probability for entry and removal of the likelihood ratio score statistic being $p = 0.05$
11 and 0.10 respectively, in line with standard practice. Height and weight were included in an
12 initial step, with the all other tested physical characteristics included in a second step,
13 regardless of whether they were significant on univariate testing. BMI and arm girth were
14 omitted as they were highly related to waist girth, which is a more valid health-related
15 measure of body composition²³. Similarly, NDI was omitted as it was a composite of the four
16 factor scores, which were more specific measures. For both the lowest and highest quartiles of
17 each physical factor, results were presented as the odds of having back pain (95% confidence
18 intervals), relative to the reference category. The strength of the multivariate predictive model
19 was estimated by Nagelkerke R^2 . Alpha was set at 0.05 for the multivariate tests.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24

RESULTS

Back pain

Back pain ever was experienced by 46.0% of the participants, back pain in the past month by 28.1%, 'chronic' (lasting more than 3 months) by 11.3% and diagnosed back pain by 11.4%. Girls had a tendency ($p < 0.1$, > 0.05) for a higher prevalence of diagnosed back pain, back pain ever and back pain in the past month (see Table 2).

Insert Table 2 about here

Physical risk factors

Descriptive statistics for physical characteristics are given in Table 2. Girls had significantly higher mean scores for BMI, back endurance, sit and reach, and the motor competence factors of Persistent Control, Kinaesthetic Integration and Bimanual Dexterity. Boys obtained significantly higher mean scores for waist girth, aerobic capacity, abdominal curl number, standing long jump, basketball throw distance, grip strength and the motor competence factor of Muscle Power. Males were also taller and heavier.

Relationships between physical risk factors and back pain

Univariate results

Tables 3 and 4 display the results of univariate analyses of physical risk factors and the four measures of back pain for boys and girls.

1 *Multivariate results in boys*

2 After multivariate logistic regression analysis, increased likelihood of back pain in the last
3 month was associated with greater aerobic capacity (OR=1.65 (95% CI: 1.10-2.46)), and
4 increased likelihood of diagnosed back pain was associated with greater waist girth (OR=2.20
5 (1.11-4.36)), and both reduced flexibility (OR=1.95 (1.06-3.58)) and greater flexibility
6 (OR=2.14 (1.17-3.90)). Lower likelihood of back pain in the past month was associated with
7 greater bimanual dexterity (OR=0.58 (0.34-0.99)). There were no other significant
8 multivariate associations between the physical risk factors and the four types of back pain. The
9 Nagelkerke R² of multivariate logistic regression models ranged from 0.019 to 0.070.

10

11 *Multivariate results in girls*

12 Increased likelihood of back pain in the past month was associated with greater abdominal
13 endurance (OR=1.56 (1.018-2.38)) and there was a trend (p<0.1) for an association with
14 reduced muscle power (OR=1.43 (0.97-2.10)). Increased likelihood of chronic back pain was
15 associated with reduced kinaesthetic integration (OR=1.72 (1.02-2.92)), and increased
16 likelihood of diagnosed back pain was associated with both reduced back endurance
17 ((OR=2.05 (1.16-3.60)) and greater back endurance (OR=2.00 (1.10-3.60)). Lower likelihood
18 of back pain ever was associated with greater leg power (OR=0.58 (0.39-0.85)). There were no
19 other significant multivariate associations between the physical risk factors and the four types
20 of back pain. The Nagelkerke R² of multivariate logistic regression models ranged from 0.019-
21 0.044.

22

23 Insert Table 3 about here

24

25 Insert Table 4 about here.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25

DISCUSSION

This study confirms that back pain is common in adolescents, with almost half having experienced back pain, 20% of whom experienced prolonged episodes. It is therefore a problem requiring attention with regard to both prevention and management.

Although our univariate analyses suggested several physical factors might be related to back pain, many of these were not significant after multivariate analysis, presumably because of competition from more strongly associated factors. Our study is the first to include a wide variety of physical risk factors in the multivariate analysis, representing an advantage over previous studies in terms of permitting a more comprehensive analysis. Accordingly, only multivariate results will be discussed below. The cross-sectional approach does not allow any assumptions about the direction of any causality, but plausible mechanisms will be discussed.

Diagnosed back pain was over twice as likely in boys with the greatest waist girth. This concurs with other studies, though these studies did not find gender differences^{19,35}. Our recent research has documented a relationship between hyperlordotic postures and increased BMI in adolescents, with an associated increased risk of LBP in this postural group³⁶. It may be that the increased risk of LBP associated with waist girth in boys is linked to altered patterns of spinal loading due to excess weight. Our lack of any body composition associations in girls resembles the findings of Kujala and colleagues¹⁹, who noted an unadjusted longitudinal association between high BMI and subsequent back pain in boys only.

This is the first study to report associations between objectively measured aerobic capacity and adolescent back pain and showed that boys with the highest aerobic capacity had a greater risk

1 of back pain in the past month. Aerobically fitter boys may have been at greater risk for back
2 pain due to more prolonged or intense activity, which might increase spinal loading beyond a
3 threshold of tissue tolerance². The association of back pain with higher aerobic capacity solely
4 in boys may relate to previous findings of a higher risk of back pain in boys involved in
5 organised sport¹⁹. However, our lack of activity measurements precludes any firm
6 conclusions.

7
8 A relationship between back pain and abdominal endurance was absent for boys. However,
9 girls with *greater* abdominal endurance had a higher risk of back pain in the past month,
10 although this risk was small. This association has not been previously reported, although
11 adolescent back pain has been related to increased trunk flexor strength²⁶. The mechanism for
12 our finding is not clear although it is known that the trunk flexors can exert significant
13 flexion/compression loading forces on the lumbar spine¹⁷. Our results conflict with the
14 previous finding that *reduced* abdominal endurance is associated with back pain in both
15 genders¹⁶, although this difference may relate to differing definitions of pain. Reasons for the
16 gender difference in our study are unknown.

17
18 Trunk extensor endurance also showed no association with pain in boys, but girls showed a U
19 shaped relationship between this variable and diagnosed back pain. Previous findings have
20 either showed no effect³¹ or have shown a relationship between low extensor endurance and
21 back pain³². This inconsistency may be because previous studies were not designed to detect
22 curvilinear relationships, and highlights the importance of such study design. The finding that
23 both deficits and excesses of back muscle performance are related to LBP is supported by
24 previous reports of different LBP subgroups presenting with excesses and deficits in back
25 muscle activity levels⁹.

1
2 For the measures of limb muscle performance, only the association between greater jump
3 distance and lower risk of back pain ever in girls was significant. Adult studies have shown an
4 analogous relationship of greater back pain with reduced leg power²⁷. Although highly
5 speculative, the association could relate to those with better lower limb muscle performance
6 making greater use of the leg than trunk muscles during vigorous lower limb activities,
7 possibly reducing spinal stress¹⁷. Equally, those with back pain may avoid activities that
8 promote leg power.

9
10 The sit and reach distance, an indication of both spinal flexion and hamstring flexibility, was
11 unrelated to back pain in girls. These findings concur with results in both genders in other
12 adolescent studies using the same test^{16,24}. However, in boys there was a U shaped relationship
13 with diagnosed back pain. Again, the lack of previous observation of such a U shaped
14 relationship may relate to the linear assumptions of previous work. It is possible that both high
15 and low levels of bending flexibility could induce increased spinal strain during normal
16 activities, high flexibility possibly placing strain on neural structures³³, and low flexibility
17 potentially inducing strain on intra-articular structures¹⁹. However lower flexibility could also
18 be a result, rather than a cause, of back pain. It should be noted that the sit and reach test does
19 not differentiate hamstring from spinal flexibility and so findings should be treated with
20 caution.

21
22 Some aspects of motor competence related to back pain, with greater Bimanual Dexterity
23 associated with less risk of back pain in the past month in boys, and poorer Kinaesthetic
24 Integration linked to a greater risk of chronic back pain in the past month in girls. Bimanual
25 Dexterity, in this context, is a measure of coordination of fine motor skills across both upper

1 limbs, and Kinaesthetic Integration refers to the ability to maintain both static and dynamic
2 balance²². This is the first adolescent study to demonstrate a relationship between aspects of
3 motor competence and back pain, and it appears to concur with findings in younger children⁷
4 and adults⁸. This finding suggests that the quality of synergist coordination may be important
5 in adolescents, in addition to quantitative factors such as strength. This is consistent with
6 theories of muscle control of the spine²⁸.

7
8 The notable gender differences in the way physical risk factors associated with back pain may
9 relate to differences in the levels and types of physical activity adopted by boys and girls¹³. It
10 may also be related to anthropometric differences, as differing heights and weights may lead to
11 differing spinal torques during the same activities. It could also relate to other structural
12 differences, and should be the focus of future research.

13
14 Although back pain was associated with various physical factors, these associations were
15 weak, with Nagelkerke R^2 of multivariate logistic regression models ranging from 0.019 to
16 0.070. This could not be attributed to missed curvilinear relationships as these were accounted
17 for in the analysis. It is unlikely that the lack of strength in the measured relationships were
18 because the physical measures failed to adequately capture key physical constructs as most
19 have been validated and widely used. Back pain is not a simple construct and the lack of
20 strong relationships may also have resulted from the measures of back pain used. However we
21 used four different measures of back pain, including a primary carer report of diagnosed back
22 pain. This suggests the weaknesses associated with self-report of back pain, such as trivial
23 cases being reported, were not the reason for limited associations.

24

1 One of the strengths of this study, compared to previous work, was the broad range of physical
2 variables included in the analysis, but it is still possible that the lack of strong relationships
3 could also be a result of some physical characteristics interacting with other risk factors not
4 examined in this study. For example poor motor competence may only be important for people
5 with high exposure to spinal loading. Similarly, certain psychosocially-defined sub-groups
6 may differ in how their back pain relates to muscle performance. Hence further studies should
7 assess interactions between physical characteristics and other psychosocial and lifestyle risk
8 factors. Similarly, the lack of any sub-grouping of back pain may have led to stronger
9 relationships between physical performance characteristics and specific types of back pain
10 remaining undetected. For example, poor back muscle endurance may only be important for
11 boys with back pain associated with motor control impairments into flexion²⁹. Finally,
12 although the age of participants was very narrowly distributed, likely variations in maturation
13 could have confounded results, and future work should consider this issue. The associations
14 seen in this study between back pain and physical factors such as body composition, aerobic
15 fitness, muscle performance, flexibility and motor competence should be viewed as
16 representing just one part of the multifactorial basis of adolescent back pain.

17

18 **CONCLUSION**

19 Physical characteristics are often regarded as important risk factors in back pain development.
20 However, although some factors were associated with adolescent back pain, these differed
21 between boys and girls, and they made only a small contribution to the logistic regression
22 models for back pain. This suggests future work should explore the interaction of multiple
23 domains of risk factors (physical, lifestyle and psychosocial) and subgroups of adolescent back
24 pain, for whom different risk factors may be important.

25

1 **KEY POINTS**

2 **Findings**

3 Aspects of fitness, motor competence and body composition were related to adolescent back
4 pain and differed between genders.

5 **Implications**

6 Whilst physical characteristics were associated with back pain in adolescents, the weak and
7 varied relationships suggests adolescent back pain should not be assumed to be the same as
8 adult back pain.

9 **Caution**

10 Lifestyle and psychosocial characteristics were not included in this study. Back pain was
11 treated as a homogeneous entity, with no analysis of subgroups.

REFERENCES

1. ACHPER. *Australian Fitness Education Award*. Richmond, SA: Australian Council for Health, Physical Education and Recreation; 1996.
2. Adams MA, Dolan P. Spine biomechanics. *Journal of Biomechanics*. 2005;38:95-102.
3. Andersen LB, Wedderkopp N, Leboeuf-Yde C. Association between back pain and physical fitness in adolescents. *Spine*. 2006;31:1740–1744.
4. Bayramoglu M, Akman MN, Cetin N, Yavuz N, Ozker R. Isokinetic measurement of trunk muscle strength in women with chronic low-back. *American Journal of Physical Medicine and Rehabilitation*. 2001;80:650-655.
5. Biering-Sorensen F. Physical measurements as risk indicators for low-back trouble over a one-year period. *Spine*. 1984;9:106-119.
6. Boreham C A, Twisk J, Murray L, Gallagher A, Savage M. Relationships between aerobic fitness, physical activity and arterial compliance in young adults. *Medicine Science Sports and Exercise*. 2003;35:S72.
7. Cardon G, De Bourdeaudhuij I, De Clercq D, Philippaerts R, Verstraete S, Geldhof E. Physical fitness, physical activity, and self-reported back and neck pain in elementary schoolchildren. *Pediatric Exercise Science*. 2004;16:147-157.
8. Cholewicki J, Silfies SP, Shah RA, Greene HS, Reeves NP, Alvi K, Goldberg B. Delayed Trunk Muscle Reflex Responses Increase the Risk of Low Back Injuries. *Spine*. 2005;30:2614-2620.
9. Dankaerts W, O’Sullivan PB, Burnett AF, Straker LM. Altered patterns of superficial trunk muscle activation during sitting in non-specific chronic low back pain patients: Importance of subclassification. *Spine*. 2006;31:2017-2023.

10. Gutin B, Basch C, Shea S, Contento I, De Lozier M, Rips J, Irigogen M, Zybert P. Blood pressure, fitness, and fatness in 5- and 6-year-old children. *Journal of the American Medical Association*. 1990;264:1123–1127.
11. Harreby, M., B. Nygaard, Jessen T, Larsen E, Storr-Paulsen A, Lindahl A, Fisher I, Laegaard E. Risk factors for low back pain in a cohort of 1389 Danish school children: An epidemiologic study. *European Spine Journal*. 1999;8:444-450.
12. Hayden JA, van Tulder MW, Malmivaara A, Koes BW. Exercise therapy for treatment of non-specific low back pain. *Cochrane Database of Systematic Reviews*. 2005. Issue 3. Art. No.: CD000335. DOI: 10.1002/14651858.CD000335.pub2.
13. Hovell MF, Sallis JF, Kolody B, McKenzie TL. Children's physical activity choices: A developmental analysis of gender, intensity levels and time. *Pediatric Exercise Science*. 1999;11:158-168.
14. Jones MA, Hitchen PJ. The prevalence of low back pain in British school children. *Journal of Sports Science*. 2000;18:15
15. Jones M, Stratton G, Reilly T, Unnithan VB. Measurement error associated with spinal mobility measures in children with and without low-back pain. *Acta Paediatrica*. 2002;91:1339–1343.
16. Jones M, Stratton G, Reilly T, Unnithan VB. Biological risk indicators for recurrent non-specific low back pain in adolescents. *British Journal of Sports Medicine*. 2005;9:137-140.
17. Kavcic N, Grenier S, McGill SM. Quantifying Tissue Loads and Spine Stability While Performing Commonly Prescribed Low Back Stabilization Exercises. *Spine*. 2004;29:2319-2329.
18. . Kujala, UM, Taimela S, Erkintalo M., Salminen JJ, Kaprio J. Low-back pain in adolescent athletes. *Medicine and Science in Sports and Exercise*. 1996;28,165-170

19. Kujala, UM, Taimela S, Oksanen A, Salminen JJ. Lumbar mobility and low back pain during adolescence. A longitudinal three-year follow-up study in athletes and controls. *American Journal of Sports Medicine*. 1997;25:363-368.
20. Markovic G, Dizdar D, Jukic I, Cardinale M. Reliability and factorial validity of squat and countermovement jump tests. *Journal of Strength and Conditioning Research*. 2004;18:551-555.
21. McArdle W, Katch F, Pechar G, Jacobson L, Ruck S. Reliability and interrelationships between maximal oxygen intake, physical work capacity and step-test scores in college women. *Medicine and Science in Sports*. 1972; 4:182-186.
22. McCarron L. *McCarron assessment of neuromotor development: Fine and gross motor abilities*. 3rd ed. Dallas TX, USA: McCarron-Dial systems inc; 1997.
23. McCarthy HD, Ellis SM, Cole TJ. Central overweight and obesity in British youth aged 11-16 years: Cross sectional surveys of waist circumference. *BMJ*. 2003;326:624-626.
24. Mikkelsen LO, Nupponen H, Kaprio J, Kautiainen H, Mikkelsen M, Kujala UM. Adolescent flexibility, endurance strength, and physical activity as predictors of adult tension neck, low back pain, and knee injury: A 25 year follow up study. *British Journal of Sports Medicine*. 2006;40:107-113.
25. Moreau CE, Green BN, Johnson CD, Moreau SR. Isometric back extension endurance tests: a review of the literature. *Journal of Manipulative and Physiological Therapeutics*. 2001;24:110-122.
26. Newcomer K, Sinaki M. Low back pain and its relationship to back strength and physical activity in children. *Acta Paediatrica*. 1996;85:1433-1439.

- 27.** Novy DM, Simmonds MJ, Olson SL, Lee CE, Jones SC. Physical performance: Differences in men and women with and without low back pain. *Archives of Physical Medicine and Rehabilitation*. 1999;80:195-198.
- 28.** O’Sullivan P. Diagnosis and classification of chronic low back pain disorders: Maladaptive movement and motor control impairments as underlying mechanism. *Manual Therapy*. 2005;10:242–255.
- 29.** O’Sullivan PB, Mitchell T, Bulich P, Waller R, Holte J. The relationship between posture and back muscle endurance in industrial workers with flexion-related low back pain. *Manual Therapy*. 2006;11: 264–271.
- 30.** Patterson P, Wiksten DL, Ray L, Flanders C, Sanphy D. The validity and reliability of the back saver sit-and-reach test in middle school girls and boys. *Research Quarterly for Exercise and Sport*. 1996;67: 448-451.
- 31.** Salminen JJ, Erkontalo M, Laine M , Pentti J. Low back pain in the young. A prospective three-year follow-up study of subjects with and without low back pain. *Spine*. 1995;20:2101-2107.
- 32.** Salminen JJ, Maki P, Oksanen A, Pentti J. Spinal mobility and trunk muscle strength in 15-year-old schoolchildren with and without low-back pain. *Spine*. 1992;17:405-411.
- 33.** Samartzis D, Lubicky J, Herman J, Kalluri P Shen F. Symptomatic Cervical Disc Herniation in a Pediatric Klippel-Feil Patient: The Risk of Neural Injury Associated With Extensive Congenitally Fused Vertebrae and a Hypermobility Segment. *Spine*. 2006; 31:E335-E338
- 34.** Shrier I, Feldman D, Klvana J, Rossignol M, Abenhaim L. Concurrent validity of abdominal endurance and abdominal strength testing in adolescents. *Medicine and Science in Sports and Exercise*. 1998;30:S214

35. Sjolie AN. Low-back pain in adolescents is associated with poor hip mobility and high body mass index. *Scandinavian Journal of Medicine and Science in Sports*. 2004;14:168-175.
36. Smith A, O'Sullivan P, Straker L. Classification of sagittal thoraco-lumbo-pelvic alignment of the adolescent spine in standing using two-dimensional photographic images. *Spine*. 2008;(in press).
37. Straker L.M, O'Sullivan PB, Smith A, Perry M. Computer use and habitual spinal posture in Australian adolescents. *Public Health Reports*. 2007;122:634-643.
38. Tsimeas PD, Tsiokanos AL, Koutedakis Y, Tsigilis N, Kellis S. Does living in urban or rural settings affect aspects of physical fitness in children? An allometric approach. *British Journal of Sports Medicine*. 2005;39:671-674.
39. Verbunt JA, Sieben JM, Seelen HA, Vlaeyen JW, Bousema EJ, van der Heijden GT, Knottnerus JA. Decline in physical activity, disability and pain related fear in sub-acute low back pain. *European Journal of Pain*. 2005;9:417-425.
40. Vikat A, Rimpela M, Salminen JJ, Rimpela A, Savolainen A, Virtanen SM. Neck or shoulder pain and low back pain in Finnish adolescents. *Scandinavian Journal of Public Health*. 2000;28:164-173.

<i>Test</i>	<i>Details</i>	<i>Competence assessed</i>
<i>Rod slide</i>	<i>Moving a bead as slowly and smoothly as possible along a rod, repeated with both hands.</i>	<i>PC</i>
<i>Finger/nose finger</i>	<i>The accuracy and smoothness of movement of the index finger from nose to opposite hand's index finger is observed, repeated on both sides. Eyes open and closed.</i>	<i>PC</i>
<i>Hand strength</i>	<i>Hand grip strength is measured with a hand dynamometer, repeated on both sides</i>	<i>MP</i>
<i>Standing long jump</i>	<i>The distance and quality of a two footed jump is recorded.</i>	<i>MP</i>
<i>Heel toe walk</i>	<i>The quality of forward and backwards walking along a 10 foot line is recorded.</i>	<i>KI</i>
<i>Standing on one leg</i>	<i>With eyes open the time the participant can balance on each leg is recorded, repeated with eyes closed</i>	<i>KI</i>
<i>Beads on rod</i>	<i>Number of cylindrical beads placed on a metal rod held in non-dominant hand in 30 seconds, repeated with eyes open and then closed</i>	<i>BD</i>
<i>Nut and bolt</i>	<i>Time taken to turn a large bolt, held in the dominant hand, fully onto a nut, repeated with a small bolt.</i>	<i>BD</i>
<i>Finger tapping</i>	<i>Number and quality of taps of index finger in 10 seconds, repeated on both hands</i>	-
<i>Beads in box</i>	<i>Number of beads moved from one box to an adjacent box in 30 seconds, repeated for both hands.</i>	-

TABLE 1. Summary of MAND tests. PC=Persistent Control, MP=Muscle Power, KI=Kinaesthetic Integration, BD=Bimanual Dexterity.

Pain variable	All Participants % (count) with history of pain	Male % (count) with history of pain	Female % (count) with history of pain			Gender difference
						P
<i>Back pain ever</i>	46.0 (734)	43.7 (356)	48.5 (378)			0.052
<i>Back pain in last month</i>	28.1 (449)	26.1 (213)	30.2 (236)			0.070
<i>Chronic back pain</i>	11.3 (180)	10.8 (88)	11.8 (92)			0.535
<i>Diagnosed back pain</i>	11.4 (177)	9.9 (79)	12.9 (98)			0.069
Physical variable	All Participants mean (sd)	Males		Females		Gender difference
		<i>mean (sd)</i>	<i>IQR</i>	<i>mean (sd)</i>	<i>IQR</i>	P
<i>Height</i>	1.64 (0.08)	1.66 (0.09)	1.61-1.73	1.62 (0.06)	1.50-1.67	<0.001
<i>Weight</i>	57.7 (13.2)	58.6 (14.1)	50.0-66.0	56.7 (12.1)	49.2-61.9	0.004
<i>BMI</i>	21.29 (4.15)	21.05 (4.14)	18.30-22.79	21.53 (4.16)	18.91-23.38	0.022
<i>Waist girth (cm)</i>	75.5 (10.8)	76.3 (11.4)	68.5-81.0	74.6 (10.1)	67.5-79.2	0.002
<i>Arm circumference (cm)</i>	25.2 (3.3)	25.3 (3.4)	23.0-27.2	25.1 (3.3)	23.0-27.0	0.244
<i>PWC₁₇₀ score (W)</i>	111.2 (29.9)	124.3 (31.7)	102.5-143.8	97.2 (19.9)	84.9-108.2	<0.001
<i>Back muscle endurance (seconds)</i>	80.9 (60.4)	77.8 (60.1)	26.0-121.0	84.2 (60.5)	32.0-120.0	0.034
<i>Abdominal muscle endurance (number of curls in 3min)</i>	21.4 (17.4)	25.4 (18.8)	12.0-36.0	17.2 (14.6)	7.0-23.0	<0.001
<i>Standing long jump distance (metres)</i>	1.46 (0.29)	1.59 (0.28)	1.42-1.78	1.32 (0.23)	1.4217-1.47	<0.001
<i>Basketball throw (metres)</i>	5.3 (1.0)	5.7 (1.0)	5.1-6.4	4.8 (0.7)	4.4-5.2	<0.001
<i>Total hand strength (kg)</i>	51.8 (13.5)	57.0 (14.8)	47.0-66.0	46.3 (9.1)	40.0-52.0	<0.001

<i>Sit and reach distance (right leg*) (cm)</i>	24.8 (9.1)	21.1 (8.0)	16.0-26.5	28.8 (8.6)	23.3-35.0	<0.001
<i>NDI score</i>	97.2 (17.4)	97.3 (18.1)	85.0-111.0	97.0 (16.6)	85.0-110.0	0.758
<i>Persistent Control factor score</i>	103.3 (25.4)	99.9 (26.4)	80.0-125.0	106.8 (23.7)	90.0-125.0	<0.001
<i>Muscle Power factor score</i>	95.9 (20.2)	102.4 (19.8)	90.0-120.0	89.2 (18.5)	75.0-100.0	<0.001
<i>Kinaesthetic Integration factor score</i>	96.9 (15.2)	96.7 (15.7)	85.0-110.0	97.2 (14.7)	90.0-110.0	<0.001
<i>Bimanual Dexterity factor score</i>	97.1 (19.3)	95.1 (19.3)	85.0-110.0	99.1 (19.1)	85.0-110.0	<0.001

TABLE 2. Summary of pain prevalence and performance in the physical tests in boys and girls. IQR=interquartile range.

*The right leg sit and reach distance did not differ from the left for boys or girls, so all analyses used the right sit and reach distance.

		% with back pain			low vs IQR		high vs IQR	
		low	IQR	high	OR	95%CI	OR	95%CI
Back Pain Ever								
Height		38.0	42.7	51.9	0.82	0.59-1.15	1.45*	1.02-2.06
Weight		36.5	44.9	48.5	0.70*	0.50-1.00#	1.16	0.83-1.62
Body composition	BMI	38.9	44.0	47.2	0.81	0.58-1.14	1.14	0.81-1.60
	Waist girth	35.4	45.7	48.0	0.65*	0.46-0.92	1.09	0.78-1.54
	Arm girth	37.7	45.7	46.8	0.72	0.51-1.00	1.04	0.74-1.47
Aerobic performance	PWC ₁₇₀	39.4	42.2	52.9	0.89	0.63-1.26	1.53*	1.08-2.17
Muscle performance	Back end	46.3	40.6	46.2	1.27	0.90-1.78	1.26	0.89-1.77
	Curls	40.8	43.2	45.9	0.91	0.64-1.28	1.11	0.79-1.57
	Jump	41.4	46.6	39.9	0.81	0.58-1.14	0.76	0.54-1.08
	Basketball	36.6	45.3	47.0	0.70*	0.49-0.99	1.07	0.77-1.51
Flexibility	Grip strength	36.2	46.4	47.4	0.66*	0.47-0.92	1.04	0.74-1.47
	Sit and reach	42.1	43.9	42.5	0.93	0.66-1.31	0.95	0.67-1.33
Motor competence	NDI	43.0	44.9	42.4	0.93	0.66-1.30	0.90	0.64-1.27
	PC	43.8	46.0	39.6	0.91	0.65-1.28	0.77	0.54-1.09
	MP	39.0	44.4	48.1	0.80	0.57-1.12	1.16	0.78-1.71
	KI	41.0	45.8	41.7	0.82	0.59-1.15	0.85	0.58-1.23
	BD	45.3	44.8	38.4	1.02	0.75-1.38	0.77	0.51-1.15
Back Pain Month								
Height		22.5	24.6	33.7	0.89	0.60-1.32	1.56*	1.07-2.28
Weight		21.2	26.1	31.2	0.76	0.51-1.14	1.28	0.89-1.86
Body composition	BMI	21.7	25.3	32.2	0.82	0.55-1.22	1.40	0.97-2.03
	Waist girth	20.6	26.8	30.1	0.71	0.47-1.06	1.18	0.81-1.72
	Arm girth	20.3	26.9	31.3	0.69	0.47-1.03	1.24	0.85-1.80
Aerobic performance	PWC ₁₇₀	21.2	24.6	35.2	0.83	0.55-1.25	1.67**	1.15-2.43
Muscle performance	Back end	25.9	26.1	24.6	0.99	0.67-1.45	0.92	0.62-1.37
	Curls	25.5	25.1	27.0	1.02	0.69-1.52	1.11	0.75-1.63
	Jump	24.2	26.7	26.6	0.88	0.59-1.30	0.99	0.67-1.47
	Basketball	22.8	25.6	29.2	0.86	0.58-1.27	1.20	0.82-1.75
Flexibility	Grip strength	19.3	28.3	29.9	0.60*	0.40-0.90	1.08	0.74-1.57
	Sit and reach	25.7	25.9	23.5	0.99	0.67-1.46	0.88	0.59-1.30
Motor competence	NDI	25.0	26.6	26.3	0.92	0.63-1.36	0.99	0.67-1.45
	PC	27.4	26.2	24.9	1.06	0.73-1.55	0.93	0.63-1.38
	MP	22.0	27.3	27.9	0.75	0.51-1.11	1.03	0.67-1.60
	KI	24.1	27.8	24.3	0.83	0.57-1.20	0.84	0.54-1.29
	BD	28.9	26.3	20.3	1.14	0.81-1.60	0.71	0.44-1.16
Back Pain Chronic								
Height		7.5	10.3	16.0	0.71	0.39-1.29	1.67*	1.01-2.77
Weight		4.4	12.2	14.4	0.33**	0.16-0.69	1.21	0.74-1.97
Body composition	BMI	8.4	11.4	12.1	0.71	0.40-1.27	1.06	0.63-1.79
	Waist girth	5.7	12.0	13.3	0.45*	0.23-0.86	1.12	0.67-1.86
	Arm girth	5.6	12.8	12.9	0.41**	0.22-0.77	1.01	0.61-1.69
Aerobic performance	PWC ₁₇₀	9.3	10.6	13.0	0.87	0.48-1.55	1.25	0.74-2.13
Muscle performance	Back end	12.2	9.7	12.3	1.29	0.76-2.20	1.31	0.76-2.24
	Curls	11.7	9.8	12.2	1.22	0.71-2.10	1.28	0.75-2.19
	Jump	7.6	12.8	10.1	0.56	0.31-1.02	0.77	0.44-1.34
	Basketball	8.9	10.5	13.4	0.84	0.47-1.50	1.32	0.79-2.22
	Grip strength	6.4	12.0	13.4	0.50*	0.27-0.93	1.13	0.68-1.89
Flexibility	Sit and reach	11.9	11.5	8.0	1.04	0.62-1.76	0.67	0.37-1.22
Motor competence	NDI	12.5	10.4	10.6	1.23	0.73-2.08	1.02	0.59-1.77
	PC	12.5	10.6	9.6	1.20	0.71-2.01	0.90	0.51-1.58
	MP	8.0	12.4	10.1	0.62	0.35-1.10	0.79	0.42-1.50
	KI	12.3	10.8	9.0	1.16	0.70-1.92	0.82	0.43-1.56
	BD	12.5	10.7	7.5	1.18	0.74-1.90	0.68	0.33-1.40
Diagnosed Back Pain								
Height		8.1	9.2	14.0	0.87	0.48-1.58	1.62	0.94-2.78
Weight		5.1	9.7	15.4	0.49	0.24-1.01	1.69*	1.01-2.81
Body composition	BMI	6.0	9.1	15.9	0.64	0.33-1.26	1.90*	1.13-3.17

<i>composition</i>	<i>Waist girth</i>	4.5	9.4	16.9	0.46*	0.22-0.96	1.96*	1.18-3.26
	<i>Arm girth</i>	4.9	10.4	14.9	0.44*	0.22-0.88	1.50	0.90-2.51
<i>Aerobic</i>	<i>PWC170</i>	10.4	9.7	11.2	1.09	0.61-1.93	1.17	0.66-2.07
<i>Muscle performance</i>	<i>Back end</i>	13.0	9.3	7.3	1.47	0.86-2.51	0.77	0.40-1.50
	<i>Curls</i>	11.2	8.1	10.6	1.43	0.81-2.54	1.35	0.75-2.43
	<i>Jump</i>	11.7	9.7	8.4	1.23	0.71-2.12	0.85	0.46-1.58
	<i>Basketball</i>	9.0	8.7	13.3	1.03	0.57-1.87	1.61	0.94-2.77
	<i>Grip strength</i>	8.3	9.3	13.4	0.89	0.49-1.61	1.52	0.88-2.61
<i>Flexibility</i>	<i>Sit and reach</i>	11.6	6.9	13.0	1.77	0.99-3.17	2.02*	1.14-3.59
<i>Motor competence</i>	<i>NDI</i>	10.1	9.0	11.1	1.14	0.64-2.03	1.26	0.71-2.24
	<i>PC</i>	10.0	10.8	7.9	0.91	0.53-1.59	0.71	0.38-1.32
	<i>MP</i>	10.6	9.7	9.0	1.10	0.64-1.91	0.93	0.46-1.85
	<i>KI</i>	11.8	10.0	6.5	1.20	0.71-2.02	0.62	0.30-1.30
	<i>BD</i>	11.0	9.1	10.2	1.23	0.74-2.05	1.13	0.57-2.22

TABLE 3. Univariate relationship between physical performance characteristics and back pain in boys

OR = Odds Ratio, 95%CI = 95% confidence interval, IQR = interquartile range, BMI = body mass index, PWC = physical work capacity, NDI = Neuromuscular Developmental Index, PC = persistent control, MP = muscle power, KI = kinesthetic integration, BD = bimanual dexterity. *P<0.05, **P<0.01. #The actual lower CI was <1.00 but rounded up to 1.00, so OR was still significantly less than 1.00.

		% with back pain			low vs IQR		high vs IQR	
		low	IQR	high	OR	95%CI	OR	95%CI
Back Pain Ever								
Height		42.3	49.9	52.4	0.74	0.57-1.11	1.11	0.78-1.57
Weight		45.2	47.4	54.1	0.91	0.13-1.31	1.31	0.93-1.85
Body composition	BMI	48.2	45.4	54.7	1.12	0.79-1.58	1.45*	1.03-2.06
	Waist girth	43.2	47.2	55.9	0.85	0.05-1.42	1.42	0.99-2.01
	Arm girth	44.9	48.0	55.1	0.88	0.13-1.33	1.33	0.92-1.92
Aerobic performance	PWC170	44.5	47.5	53.0	0.89	0.22-1.25	1.25	0.87-1.79
Muscle performance	Back end	49.7	46.4	49.2	1.14	0.81-1.61	1.12	0.79-1.58
	Curls	49.3	47.2	48.6	1.08	0.77-1.53	1.06	0.74-1.51
	Jump	52.7	51.5	36.7	1.05	0.00-0.55	0.55**	0.38-0.78
	Basketball	51.0	48.2	47.5	1.12	0.89-0.97	0.97	0.68-1.40
	Grip strength	49.1	47.2	49.7	1.08	0.77-1.51	1.11	0.78-1.57
Flexibility	Sit and reach	51.3	46.2	50.3	1.23	0.87-1.73	1.18	0.82-1.68
Motor competence	NDI	51.7	48.7	42.9	1.13	0.19-0.79	0.79	0.56-1.12
	PC	51.1	46.7	49.6	1.19	0.87-1.65	1.12	0.75-1.68
	MP	53.3	48.0	41.5	1.24	0.15-0.77	0.77	0.54-1.10
	KI	50.3	48.9	44.7	1.06	0.40-0.85	0.84	0.57-1.25
	BD	46.6	49.9	47.8	0.88	0.63-1.22	0.92	0.64-1.32
Back Pain Month								
Height		31.7	29.0	31.4	1.14	0.79-1.65	1.12	0.77-1.64
Weight		28.9	28.5	35.1	1.02	0.70-1.49	1.36	0.94-1.96
Body composition	BMI	31.4	26.3	36.8	1.29	0.88-1.88	1.63**	1.13-2.36
	Waist girth	26.4	30.1	34.2	0.84	0.31-1.21	1.21	0.84-1.76
	Arm girth	29.2	29.7	32.9	0.98	0.68-1.40	1.16	0.78-1.72
Aerobic performance	PWC170	28.6	29.2	33.7	0.97	0.28-1.24	1.23	0.84-1.81
Muscle performance	Back end	34.9	27.3	28.3	1.43	0.98-2.07	1.05	0.71-1.55
	Curls	33.7	26.3	31.5	1.43	0.98-2.07	1.29	0.88-1.90
	Jump	32.7	33.2	21.3	0.98	0.68-1.41	0.55**	0.36-0.82
	Basketball	33.9	29.0	28.6	1.26	0.87-1.81	0.98	0.66-1.47
	Grip strength	32.9	28.2	30.5	1.25	0.87-1.79	1.12	0.76-1.64
Flexibility	Sit and reach	30.4	29.6	29.5	1.04	0.72-1.51	1.00	0.67-1.47
Motor competence	NDI	31.7	30.5	27.4	1.06	0.44-0.86	0.86	0.58-1.27
	PC	33.2	29.3	26.6	1.20	0.56-0.87	0.87	0.56-1.37
	MP	36.3	29.5	23.3	1.36	0.13-0.73	0.73	0.48-1.10
	KI	29.4	31.9	25.0	0.89	0.62-1.28	0.71	0.46-1.10
	BD	29.5	31.0	28.6	0.93	0.65-1.33	0.89	0.60-1.32
Back Pain Chronic								
Height		10.1	13.2	10.6	0.74	0.43-1.27	0.79	0.45-1.36
Weight		12.7	11.3	11.9	1.14	0.68-1.93	1.06	0.62-1.81
Body composition	BMI	13.9	9.8	13.5	1.49	0.88-1.52	1.43	0.84-2.44
	Waist girth	11.9	11.1	13.4	1.14	0.68-1.93	1.06	0.62-1.81
	Arm girth	11.9	10.6	14.6	1.15	0.69-1.92	1.44	0.83-2.49
Aerobic performance	PWC170	10.4	11.1	14.9	0.93	0.21-1.40	1.40	0.83-2.37
Muscle performance	Back end	13.3	10.0	11.0	1.39	0.82-2.36	1.12	0.64-1.96
	Curls	12.1	11.3	9.9	1.08	0.63-0.87	0.87	0.49-1.55
	Jump	13.5	11.0	10.1	1.26	0.76-0.91	0.91	0.51-1.62
	Basketball	12.8	10.5	13.0	1.26	0.74-2.13	1.28	0.74-2.24
	Grip strength	14.6	9.0	12.8	1.71*	1.02-2.88	1.48	0.85-2.58
Flexibility	Sit and reach	11.3	11.3	12.1	1.01	0.59-1.73	1.09	0.63-1.89
Motor competence	NDI	11.9	10.4	12.6	1.16	0.68-1.99	1.24	0.72-2.13
	PC	10.0	11.4	14.5	0.87	0.34-1.33	1.33	0.74-2.38
	MP	14.4	9.0	12.5	1.70*	1.01-2.85	1.44	0.81-2.54
	KI	16.0	10.4	9.1	1.63*	1.00-2.67##	0.86	0.44-1.67
	BD	12.7	10.6	11.5	1.22	0.73-2.03	1.10	0.62-1.93
Diagnosed Back Pain								
Height		9.5	14.7	12.7	0.61	0.35-1.05	0.84	0.50-1.42
Weight		10.5	13.7	13.6	0.74	0.43-1.27	0.99	0.60-1.65
Body composition	BMI	10.6	13.5	13.2	0.76	0.44-1.32	0.98	0.59-1.63
	Waist girth	12.5	13.1	12.6	0.95	0.56-1.61	0.96	0.56-1.62

	<i>Arm girth</i>	10.9	13.3	14.7	0.80	0.48-1.34	1.13	0.67-1.93
<i>Aerobic</i>	<i>PWC170</i>	11.4	12.6	15.3	0.89	0.51-1.57	1.25	0.75-2.10
<i>Muscle</i>	<i>Back end</i>	15.0	9.5	15.1	1.69*	1.00-2.87##	1.71*	1.00-2.91##
<i>performance</i>	<i>Curls</i>	12.8	12.6	10.4	1.01	0.60-1.70	0.80	0.45-1.43
	<i>Jump</i>	11.9	12.8	13.7	0.92	0.54-1.56	1.07	0.64-1.81
	<i>Basketball</i>	10.0	14.0	12.7	0.68	0.39-1.19	0.90	0.52-1.55
	<i>Grip strength</i>	10.1	13.2	15.2	0.75	0.43-1.28	1.19	0.72-1.96
<i>Flexibility</i>	<i>Sit and reach</i>	13.5	12.4	11.5	1.10	0.66-1.84	0.91	0.52-1.61
<i>Motor</i>	<i>NDI</i>	11.11	11.81	16.30	0.93	0.54-1.61	1.45	0.88-2.41
<i>competence</i>	<i>PC</i>	12.05	12.35	15.97	0.97	0.59-1.60	1.35	0.76-2.39
	<i>MP</i>	11.43	12.84	14.04	0.88	0.52-1.48	1.11	0.65-1.88
	<i>KI</i>	12.50	13.04	13.28	0.95	0.57-1.59	1.02	0.57-1.83
	<i>BD</i>	10.92	13.88	13.14	0.76	0.46-1.27	0.94	0.55-1.60

TABLE 4. Univariate relationship between physical performance characteristics and back pain in girls.

OR = Odds Ratio, 95%CI = 95% confidence interval, IQR = interquartile range, BMI = body mass index, PWC = physical work capacity, NDI = Neuromuscular Developmental Index, PC = persistent control, MP = muscle power, KI = kinesthetic integration, BD = bimanual dexterity. *P<0.05, **P<0.01. ##The actual lower CI was >1.00 but rounded down to 1.00, so OR was still significantly greater than 1.00.