

2017

Trends in chlamydia and gonorrhoea testing and positivity in Western Australian Aboriginal and non-Aboriginal women 2001-2013: a population based cohort study

J. Reekie

B. Donovan

R. Guy

J. Hocking

J. Kaldor

See next page for additional authors

Follow this and additional works at: https://researchonline.nd.edu.au/med_article



Part of the [Medicine and Health Sciences Commons](#)

This article was originally published as:

Reekie, J., Donovan, B., Guy, R., Hocking, J., Kaldor, J., Mak, D., Pearson, S., Preen, D., Wand, H., Ward, J., & Liu, B. (2017). Trends in chlamydia and gonorrhoea testing and positivity in Western Australian Aboriginal and non-Aboriginal women 2001-2013: a population based cohort study. *Sexual Health*, 14 (6), 574-580.

<http://doi.org/10.1071/SH16207>

This article is posted on ResearchOnline@ND at https://researchonline.nd.edu.au/med_article/870. For more information, please contact researchonline@nd.edu.au.



Authors

J. Reekie, B. Donovan, R. Guy, J. Hocking, J. Kaldor, D. Mak, S. Pearson, D. Preen, H. Wand, J. Ward, and B. Liu

This is the author's version of the article published as:

Reeje, J., Donovan, B., Guy, R., hocking, J., Kaldor, J., Mak, D. ... Liu, B. (2017). Trends in chlamydia and gonorrhoea testing and positivity in Western Australian Aboriginal and non-Aboriginal women 2001-2013: a population based cohort study. *Sexual Health*, 14(6), 574-580. Doi: 10.1071/SH16207

1 **Trends in chlamydia and gonorrhoea testing and positivity in Western Australian Aboriginal and**
2 **non-Aboriginal women 2001-2013: a population based cohort study**

3 Joanne Reekie^{A*}, Basil Donovan^{A,B}, Rebecca Guy^A, Jane S Hocking^C, John M Kaldor^A, Donna B Mak^D,
4 Sallie Pearson^E, David Preen^F, Handan Wand^A, James Ward^G and Bette Liu^H on behalf of the chlamydia
5 and reproductive health outcome investigators

6

7 ^AKirby Institute, UNSW Australia, Sydney, Australia; ^BSydney Sexual Health Centre, Sydney Hospital,
8 Sydney, Australia; ^CSchool of Population and Global Health, University of Melbourne, Melbourne,
9 Australia; ^DSchool of Medicine, The University of Notre Dame, Fremantle Australia; ^EFaculty of
10 Pharmacy and School of Public Health, University of Sydney, Sydney, Australia ;^FCentre for Health
11 Services Research, University of Western Australia, Perth, Australia; ^GSouth Australian Health and
12 Medical Research Institute, Adelaide, Australia; ^HSchool of Public Health and Community Medicine,
13 UNSW Australia, Sydney, Australia

14

15 Word Count: 2699

16 Abstract: 242

17

18

19 *Corresponding author:

20 Joanne Reekie

21 Email: jreekie@kirby.unsw.edu.au

22

23 **Abstract**

24 **Aims:** To examine trends in chlamydia and gonorrhoea testing and positivity in Aboriginal and non-
25 Aboriginal women of reproductive age.

26 **Methods:** A cohort of 318002 women, born between 1974-1995, residing in Western Australia (WA)
27 was determined from birth registrations and the 2014 electoral roll. This cohort was then
28 probabilistically linked to all records of chlamydia and gonorrhoea nucleic acid amplification tests
29 (NAAT) conducted between 1st January 2001 and 31st December 2013 by two large WA pathology
30 laboratories. Trends in chlamydia and gonorrhoea testing and positivity were investigated over time
31 and stratified by Aboriginality and age group.

32 **Results:** The proportion of women tested annually for chlamydia increased significantly between 2001
33 and 2013 from 24% to 37% in Aboriginal and 4.0% to 8.5% in non-Aboriginal women (both p-values
34 <0.001). Concurrent testing was high (>80%) and so patterns of gonorrhoea testing were similar.
35 Chlamydia and gonorrhoea positivity were substantially higher in Aboriginal compared to non-
36 Aboriginal women; age-, region- and year-adjusted Incidence Rate Ratio's 1.52(95%CI 1.50-1.69,
37 p<0.001), and 11.80(95%CI 10.77-12.91, p<0.001) respectively. Chlamydia positivity increased
38 significantly in non-Aboriginal women aged 15-19 peaking in 2011 at 13.3%(12.5%-14.2%); trends
39 were less consistent among 15-19 year old Aboriginal women but positivity also peaked in 2011 at
40 18.5%(16.9%-20.2%). Gonorrhoea positivity was 9.7%(9.3%-10.1%), 6.7%(6.4%-7.0%), 4.7%(4.4%-
41 5.0%), and 3.1%(2.8%-3.4%) among Aboriginal women aged respectively 15-19, 20-24, 25-29 and ≥30
42 year, compared to <1% in all age groups in non-Aboriginal women. Over time, gonorrhoea positivity
43 declined in all age groups among Aboriginal and non-Aboriginal women.

44 **Conclusion:** Between 2001 and 2013 in WA chlamydia and gonorrhoea positivity remained highest in
45 young Aboriginal women despite chlamydia positivity increasing among young non-Aboriginal women.

46 More effective prevention strategies, particularly in young Aboriginal women are needed to address
47 these disparities.

48

49

50 **Introduction**

51 Genital *chlamydia trachomatis* infection (chlamydia) is the most frequently reported notifiable
52 infection in Australia and rates have increased substantially over the last 15 years (1-3). Notification
53 rates of *Neisseria gonorrhoea* infection (gonorrhoea) have also increased, albeit not as
54 dramatically(3). Parallel to these increases there has been an increase in testing (3).

55 Within Australia the rates of diagnosis of both chlamydia and gonorrhoea are significantly higher
56 among the Aboriginal and Torres Strait Islander population (hereafter referred to as Aboriginal)(3).
57 Further, among Aboriginal people the prevalence of chlamydia and gonorrhoea has been found to
58 vary. Chlamydia prevalence has been found to be highest in young Aboriginal people, Aboriginal
59 people living in regional areas, and Aboriginal pregnant females while the highest prevalence of
60 gonorrhoea has been reported among young Aboriginal people, and Aboriginal people living in remote
61 areas of Australia(4).

62 Annual screening for chlamydia infection has been recommended for all sexually active people aged
63 15–25 years since 2008 and for 15-29 year-olds from 2012, particularly if they are under age 20 years
64 or Aboriginal (5). However, testing for gonorrhoea is only recommended for individuals thought to be
65 at an increased risk(5). This study aimed to examine trends over 12 years (2001-2013) in chlamydia
66 and gonorrhoea testing and positivity based on individual pathology data in a cohort of Aboriginal and
67 non-Aboriginal women of reproductive age in Western Australia (WA).

68

69

70

71 **Methods**

72 *Study population*

73 This study was conducted using population-based record linkage. The linkage was conducted,
74 independent of the study investigators, by the Western Australian Data Linkage Branch (DLB) using
75 personal identifiers such as name, date of birth, address, and sex, to probabilistically link records(6).
76 Linkage accuracy using this process is high with an error rate estimated at 0.11%(7)

77 A cohort of reproductive aged women residing in WA was constructed by probabilistically linking the
78 WA Birth Registrations Data Collection, which contains a record of all children born and registered in
79 WA from 1974 onwards, and the 2014 WA Electoral Roll. Eligible women were those with either a birth
80 registration between 1974 and 1995 or a record on the WA Electoral Roll with year of birth between
81 1974 and 1995.

82 This cohort was then probabilistically linked to all records of chlamydia and gonorrhoea nucleic acid
83 amplification tests (NAAT) conducted between 1st January 2001 and 31st December 2013 by two large
84 WA pathology laboratories. The type of test (chlamydia or gonorrhoea), date of referral, and test result
85 (positive, negative, or equivocal/undetermined) were supplied to the research team.

86 *Statistical Analysis*

87 Only tests recorded after the women's 15th birthday were included in the analysis. Multiple tests for
88 the same infection on the same referral date were counted as one test and considered positive if any
89 of the results were positive. A concurrent or duplex test was defined as having been tested for both
90 chlamydia and gonorrhoea on the same referral date.

91 The proportion of women tested annually at either of the two laboratories was calculated as a
92 proportion of the total number of women in the cohort who were aged ≥ 15 years old in that year. Chi-

93 squared tests were used to compare this proportion in different years (2001 to 2013) and in different
94 age-groups (15-19, 20-24, 25-29 and ≥ 30 years).

95 Positivity was defined as the number of positive tests divided by the total number of tests after
96 excluding equivocal results. The numerator and denominator could contain multiple tests (e.g. tests
97 of cure) for the same individual if that person was tested more than once during our study period.

98 Poisson regression using generalized estimating equations and robust standard errors, was used to
99 investigate trends in chlamydia and gonorrhoea positivity over time. All analyses were either adjusted
100 for or stratified by Aboriginality, age at the time of testing, and geographical region of residence.
101 Aboriginality was determined from the Indigenous Status Flag created by the DLB(8). Geographical
102 region was defined, using postcodes, according to WA Health administrative regions as; Metro: North
103 and South Metropolitan, Rural: Great Southern, Wheatbelt and South West, and Remote: Midwest,
104 Kimberly, Pilbara and Goldfields

105 *Sensitivity analysis*

106 To estimate what proportion of women may have been tested for chlamydia or gonorrhoea in labs
107 other than those we linked to(9) and whether it is likely to have changed during our study period, the
108 WA Notifiable Infectious Diseases Database (WANIDD), which contains a record of all chlamydia and
109 gonorrhoea notifications reported to the WA Department of Health under statute(10), was also linked
110 to the cohort. The proportion of chlamydia or gonorrhoea notifications with corresponding positive
111 pathology records were compared by year, age group, region and Aboriginality. Further, as women
112 may not have been resident in WA for the whole study period, analyses were repeated for the sub-
113 group of women with both a birth registration and Electoral Roll record in 2014.

114 Finally, to investigate if changes in the frequency of retesting could potentially bias the positivity
115 trends, analyses were repeated with positivity calculated using only the first positive test from a

116 woman in each year and then using only the first test ever recorded for a woman regardless of year.

117 Analyses were performed in SAS 9.4 (SAS Institute, Cary NC, USA).

118 *Ethics*

119 The study was approved by the WA Department of Health HREC (Ref #2012/73) and the WA Aboriginal

120 Health Ethics Committee (Ref 470).

121 **Results**

122 A total of 318002 women were included in the cohort; 14791 (4.7%) were Aboriginal.

123

124 *Testing trends, 2001- 2013*

125 Between 2001 and 2013, 134980 (42%) women had at least one chlamydia NAAT and 124909 (39%)
126 at least one gonorrhoea NAAT at either of the two laboratories. Testing for both chlamydia and
127 gonorrhoea was higher among Aboriginal women; 80% and 79% of Aboriginal women had been tested
128 at least once for chlamydia and gonorrhoea respectively compared to 41% and 37% of non-Aboriginal
129 women (both $p < 0.001$).

130 Between 2001 and 2013, an increase in the proportion of women tested annually for chlamydia and
131 gonorrhoea was observed, across all age-groups and in both Aboriginal and non-Aboriginal women
132 (Table 1). Among Aboriginal women aged 20-24 years old, the proportion tested annually for
133 chlamydia increased from 27.8% (95%CI 26.2%-29.4%) in 2001 to 42.5% (41.0%-44.1%) in 2013
134 ($p < 0.001$) and in non-Aboriginal women of the same age it increased from 5.1% (4.9%-5.3%) to 11.0%
135 (10.8%-11.2%) ($p < 0.001$).

136 Concurrent testing rates were $>80\%$ in both Aboriginal and non-Aboriginal women. Among Aboriginal
137 women the proportion of concurrent gonorrhoea tests, was consistently around 98% throughout the
138 study period; among non-Aboriginal women the proportion increased from 83% in 2001 to 89% in
139 2013 ($p < 0.001$). Due to high concurrent testing, the trends in gonorrhoea NAAT between 2001 and
140 2013 were similar to those of chlamydia (Table 1).

141

142 *Chlamydia positivity, 2001-2013*

143 Chlamydia positivity in the cohort was 6.8% (95%CI 6.7%-6.8%) overall although this varied with year,
144 age and Aboriginality (Figure 1). Among Aboriginal women chlamydia positivity was 16.2% (15.7%-
145 16.7%) in 15-19 year olds compared to, 9.4% (9.1%-9.8%), 5.6% (5.3%-5.9%) and 3.1% (2.8%-3.5%) in
146 those aged 20-24, 25-29 and ≥ 30 years respectively. Among non-Aboriginal women, chlamydia
147 positivity was 10.2% (10.0%-10.5%) 7.3% (7.1%-7.4%), 3.4% (3.3%-3.5%) and 1.8% (1.7%-1.9%) for
148 these age groups respectively. The results of the Poisson regression analysis (Table 2) found that after
149 adjusting for age, region, and year of test, chlamydia positivity was significantly higher in Aboriginal
150 women compared to their non-Aboriginal counterparts (adjusted Incidence Rate Ratio [aIRR] 1.56,
151 95%CI 1.50-1.69, $p < 0.001$).

152 Figure 1 shows the trends in chlamydia positivity stratified by age-group and Aboriginality. For 15-19
153 year old Aboriginal women, chlamydia positivity was relatively stable from 2001-2013 (IRR 1.00[0.99-
154 1.01] $p = 0.84$). Among Aboriginal women aged 20-24, 25-29, and ≥ 30 years there was a small but
155 significant decline in chlamydia positivity (IRR 0.99[0.98-0.99] $p = 0.03$; 0.98[0.97-1.00] $p = 0.02$; and IRR
156 0.95[0.91-0.99] $p = 0.02$ respectively).

157 Chlamydia positivity in young non-Aboriginal women increased significantly over time peaking in 2011
158 at 13.3% (12.5%-14.2%) and 8.7% (8.2%-9.3%) in women aged respectively 15-19 and 20-24 years.
159 Overall chlamydia positivity in non-Aboriginal women aged 15-19 years increased by 5% per year (IRR
160 1.05, 95%CI 1.04-1.06, $p < 0.001$) and in those aged 20-24 years by 3% per year (IRR 1.03[1.02-1.04]
161 $p < 0.001$). In non-Aboriginal women aged 25-29 years chlamydia positivity during 2001-2013 was
162 stable (IRR 1.00[0.99-1.01] $p = 0.88$) and in those ≥ 30 years, there was a significant decline in chlamydia
163 positivity (IRR 0.94[0.91-0.97] $p < 0.001$).

164

165 *Gonorrhoea positivity 2001- 2013*

166 Overall gonorrhoea positivity was 1.9% (95%CI 1.9%-2.0%) in the cohort and this varied with year, age
167 and Aboriginality (Figure 2). Gonorrhoea positivity at all ages was substantially higher in Aboriginal
168 than non-Aboriginal women, 9.7% (9.3%-10.1%), 6.7% (6.4%-7.0%), 4.7% (4.4%-5.0%), and 3.1% (2.8%-
169 3.4%) respectively in Aboriginal women aged 15-19, 20-24, 25-29 and ≥ 30 years compared to 0.6%
170 (0.5%-0.6%), 0.4% (0.3%-0.4%), 0.3% (0.3%-0.4%) and 0.2% (0.2%-0.3%) respectively among non-
171 Aboriginal women. After adjustment for age, region and year of test, gonorrhoea positivity remained
172 a significantly among Aboriginal women compared to their non-Aboriginal counterparts (aIRR
173 11.57[10.57-12.66] $p < 0.001$) (table 2). Further, compared to metropolitan WA, those residing in rural
174 WA had lower gonorrhoea positivity (aIRR 0.66[0.55-0.78] $p < 0.001$) and in remote WA significantly
175 higher gonorrhoea positivity (aIRR 1.61[1.49-1.74] $p < 0.001$).

176 Figure 2 shows the trends in gonorrhoea positivity stratified by age-group and Aboriginality. Over time
177 significant decreases in gonorrhoea positivity were observed in Aboriginal women across all age-
178 groups (all p -values < 0.001) although it remained substantially higher than in non-Aboriginal women.
179 Similarly, a significant decreases in gonorrhoea positivity were observed in non-Aboriginal women
180 aged 15-19 years (IRR 0.90 95%CI 0.87-0.94, $p < 0.001$), 20-24 years (IRR 0.93[0.90-0.97] $p < 0.001$) and
181 25-29 years (IRR 0.93[0.88-0.98] $p = 0.005$). Although no significant decline in gonorrhoea positivity was
182 observed in the oldest (≥ 30 years) age group (IRR 1.04[0.93-1.17] $p = 0.49$) this group still had the
183 lowest percentage positivity in 2013 at 0.2%.

184

185 *Sensitivity analysis*

186 Comparing notifications in WANIDD to positive pathology records, 52% (11323) of chlamydia and 87%
187 (5020) of gonorrhoea notifications that linked to women in our cohort had a corresponding positive
188 pathology test in the same year. There was no significant difference in the proportion of Aboriginal
189 and non-Aboriginal women with corresponding pathology records ($p = 0.15$ and $p = 0.14$ for chlamydia

190 and gonorrhoea respectively). However, there was a decrease, in the proportion of notifications with
191 a corresponding pathology record over time from 64% to 43% of chlamydia notifications $p<0.001$ and
192 from 88% to 83% of gonorrhoea notifications $p<0.001$ between 2001 and 2013). This trend was
193 consistent across age groups, Aboriginality and geographical region.

194 Restricting analyses to the 61% of women with a record in both the birth registrations and 2014
195 Electoral Roll was consistent with the main analysis.

196 Overall trends were mostly similar when analyses were repeated with positivity calculated using only
197 the first positive test from a woman in a particular year and excluding all subsequent tests in that year
198 (S1 Appendix Figures A and B), and also from analyses using only the first test ever recorded for a
199 woman (S1 Appendix Figures C and D). There was however a small increase in chlamydia positivity
200 found for Aboriginal women aged 15-24 years when only the first test ever recorded was used (see
201 Appendix for full results).

202

203

204 Discussion

205 In this large study of WA women we found significant increases in both chlamydia and gonorrhoea
206 testing using NAATs between 2001 and 2013. During this same period, chlamydia and gonorrhoea
207 positivity remained highest in young Aboriginal women, despite chlamydia positivity increasing among
208 young non-Aboriginal women.

209 To our knowledge this is the first study using individual pathology data to examine patterns of testing
210 and positivity over 12-years and to be able to compare these trends in Aboriginal and non-Aboriginal
211 women. In comparison, other studies have relied on data from the national notification scheme or
212 Medicare(3), been restricted to individual clinics or sexual health services, or have had substantially
213 fewer years of follow-up(9).

214 Both annual chlamydia and gonorrhoea testing rates were found to have increased during our study
215 period. While the percentage of Aboriginal women tested for chlamydia was significantly higher than
216 non-Aboriginal women in all age groups and across all years, the relative increase in chlamydia testing
217 over time was greater in non-Aboriginal compared to Aboriginal women, rising about 2-fold in non-
218 Aboriginal women compared to about 1.5-fold in Aboriginal women. Modelling work looking at the
219 benefits of routine annual chlamydia screening predicted screening ~30% of 15–24-year-old males
220 and females each year would reduce chlamydia prevalence among women by >70%(11). Assuming
221 that the 52% of notifications that linked to a positive pathology test reflects the proportion of
222 chlamydia tests undertaken at these two laboratories, testing rates among women in this age group
223 in the most recent years would be estimated to be only around 25%.

224 Consistent with data from the 2015 National Annual Surveillance Report and a report by the WA
225 Department of Health, our results showed chlamydia positivity peaking in 2011(3, 9) and young
226 Aboriginal women having the greatest risk of a positive chlamydia test(3, 12, 13). Also similar to the
227 national report, gonorrhoea positivity was markedly higher (at least 10 times greater) in young

228 Aboriginal women than non-Aboriginal women(3). Findings from the Australian Collaboration for
229 Coordinated Enhanced Sentinel Surveillance (ACCESS) project reported a significant increase in
230 chlamydia positivity among Aboriginal women (2006-2011)(13). Although we found chlamydia
231 positivity rates in this population were relatively stable over our entire study period, when the data
232 were considered over a similar time period (2007-2011) a small increase in positivity was observed,
233 which was also seen in the sensitivity analysis. A recent systematic review of the prevalence of
234 chlamydia and gonorrhoea in Aboriginal Australians reported pooled prevalence's of 12.7% (10.2%-
235 15.2%) and 10.7% (8.4%-13.0%) respectively in women(14), which although not stratified by age, were
236 similar to the positivity rates observed among 15-24 year olds in this study.

237 The introduction of NAAT in the late 1990s (enabling patient self-collected specimens), public health
238 programs to increase awareness of chlamydia, and changes to the guidelines for chlamydia testing to
239 recommend annual opportunistic screening for chlamydia infection in all sexually active young people
240 aged<30 years(5), likely explain the increases in testing for chlamydia that we observed. During 2010–
241 13, there was a scaling up of programs to increase STI testing among priority populations, particularly
242 among Aboriginalpeople(15) and this may be contributing to the observed decrease in chlamydia
243 positivity since 2011. While gonorrhoea testing is only recommend in those thought to be at high risk
244 or in areas of high prevalence(5), we found that concurrent testing of gonorrhoea with chlamydia
245 occurred throughout most of the study period. As others have suggested(16), due to increases in
246 testing, changes in gonorrhoea positivity need to be interpreted with caution. If the increased testing
247 was in low risk women, this may explain the decrease in gonorrhoea positivity that we observed,
248 rather than there being a true decrease in prevalence.

249 Despite the small decrease in recent years, the high levels of chlamydia positivity in young Aboriginal
250 and non-Aboriginal Australian women are concerning. Variations in risk behaviour, such as increasing
251 numbers of sexual partners (17, 18), which have been associated with an increased risk of chlamydia
252 (19) is one possible explanation for the changes in chlamydia positivity. The data used in this study

253 did not contain information on sexual history or behaviour so we were unable to adjust for this.
254 However, increasing chlamydia positivity has been reported in studies that were able to account for
255 changes in sexual risk factors (20). It is also possible that the increases in chlamydia positivity
256 demonstrate better targeting of testing to those at greatest risk, which could also explain the fall since
257 2011.

258 Given the large geographic distances in WA, and the differences in chlamydia and gonorrhoea
259 positivity by age group, Aboriginality and region of residence (Table 2), our analyses cannot distinguish
260 between whether the overall trends observed are indicative of a consistent pattern among all women
261 in the cohort or the result of multiple outbreaks occurring in different populations at different times.
262 A better understanding could be gained by examining trends by smaller geographic units and
263 separated by age and Aboriginality. Furthermore studies looking at differences in the structure of
264 sexual networks between Aboriginal and non-Aboriginal populations may help to provide a greater
265 understanding of the different patterns of chlamydia and gonorrhoea observed and in doing so
266 provide better insight into how to effectively address these differences.

267 Our study strengths include the use of a large representative population cohort of women with a
268 proportion identified as Aboriginal (4.7%) that is consistent with population census data for this age
269 group (4.2% in 2008)(21). Our major limitation is that we were only able to link to records from two
270 of the laboratories servicing WA, so we would not have comprehensive state-wide records for all
271 chlamydia and gonorrhoea tests. Additionally, our analyses focused on trends in NAAT testing and
272 there is the possibility that some women may have been tested by culture either in conjunction with
273 or in place of NAAT (22, 23). Therefore the numbers of tests reported is likely to be an
274 underestimation. However, the coverage from our two pathology labs, based on notifications, was
275 relatively consistent across age groups, Aboriginality, and geographical region.

276 To conclude, this study found that chlamydia positivity remained highest in young Aboriginal women,
277 at around 15%, with little change observed between 2001 and 2013, despite increases in positivity in

278 young non-Aboriginal women during the same period. Further gonorrhoea positivity was at least 10
279 times greater in young Aboriginal women than their non-Aboriginal counterparts. More effective
280 prevention strategies and continued surveillance of chlamydia and gonorrhoea testing, positivity by
281 age and risk groups are needed to address these disparities.

282

283 **Conflicts of interest**

284 None declared.

285

286 **Funding**

287 This study was funded by the National Health and Medical Research Council (NHMRC) grant

288 #APP1020628. BL, JH, BD, CR, JW, RG, JK are supported by NHMRC fellowships.

289

290 **Acknowledgements**

291 The Authors wish to thank the staff at the Western Australian Data Linkage Branch, WA Birth

292 Registrations, WA Electoral Roll, WA Department of Health and the two pathology laboratories who

293 provided data for this project. Chlamydia and Reproductive Health Outcome Investigators: B Liu, D

294 Preen, J Hocking, B Donovan, C Roberts, J Ward, D Mak, R Guy, J Kaldor, S Pearson, L Stewart, H Wand

295 **Table 1 Proportion of women tested annually for chlamydia and gonorrhoea between 2001 and 2013, by year, age group and Aboriginality**

Percentage of women tested annually for chlamydia at either pathology laboratory								
Year	Non-Aboriginal				Aboriginal			
	15-19yrs	20-24yrs	25-29yrs	≥30yrs*	15-19yrs	20-24yrs	25-29yrs	≥30yrs*
2001	3.5	5.1	3.2		23.4	27.8	20.6	
2002	3.6	5.3	3.4		24.5	29.4	24.4	
2003	3.8	5.7	3.6		25.7	30.9	25.4	
2004	4.0	5.9	4.1	1.7	27.2	31.8	29.3	19.1
2005	5.2	8.0	5.5	3.3	29.3	32.6	28.8	21.2
2006	6.0	8.9	6.3	4.1	30.0	32.8	28.7	23.8
2007	6.9	10.2	7.5	5.0	32.5	36.7	31.6	24.7
2008	6.9	10.5	9.0	6.2	32.3	37.6	33.7	24.4
2009	7.0	10.7	9.2	6.2	32.1	38.5	32.8	26.2
2010	7.0	10.6	9.2	6.1	32.8	38.3	32.5	25.4
2011	8.7	10.6	9.9	6.2	37.8	39.8	33.8	25.2
2012	11.1	11.5	9.9	6.1	45.7	40.2	37.0	28.2
2013	12.2	11.0	9.9	6.0	51.2	42.5	38.3	27.7
Percentage of women tested annually for gonorrhoea at either pathology laboratory								
Year	Non-Aboriginal				Aboriginal			
	15-19yrs	20-24yrs	25-29yrs	≥30yrs*	15-19yrs	20-24yrs	25-29yrs	≥30yrs*
2001	3.0	4.3	2.7		23.2	27.7	20.0	
2002	3.2	4.6	2.9		24.4	29.3	24.0	
2003	3.4	5.0	3.1		25.4	30.4	25.1	
2004	3.5	5.0	3.5	1.5	27.0	31.6	28.8	18.7
2005	4.5	6.9	4.8	2.8	29.0	32.2	28.5	21.1
2006	5.2	7.7	5.5	3.5	29.4	32.3	28.1	23.2
2007	6.1	9.2	6.6	4.3	32.3	36.2	31.1	24.4
2008	6.4	9.5	7.9	5.3	31.7	36.7	32.6	24.0
2009	6.4	9.6	8.0	5.2	30.9	37.2	31.9	25.9
2010	6.4	9.5	8.0	5.2	32.2	37.5	31.7	24.9
2011	8.0	9.5	8.5	5.2	36.9	38.9	32.8	24.5
2012	10.4	10.5	8.7	5.3	45.2	39.6	36.4	27.8
2013	11.4	11.0	8.8	5.2	50.4	42.1	38.0	27.4

296 *Trends for the oldest age group (≥ 30 years) are reported from 2004, as per our inclusion criteria the earliest date of birth for women included in the

297 cohort was 1st January 1974.

298

299 **Table 2 Factors associated with chlamydia and gonorrhoea positivity**

	Univariate		Multivariate	
Factors associated with chlamydia positivity				
	IRR	P-value	aIRR	P-value
Age group				
15-19	1.56 (1.51-1.61)	<.0001	1.54 (1.49-1.58)	<.0001
20-24	1 (Ref)		1 (Ref)	
25-29	0.49 (0.47-0.51)	<.0001	0.50 (0.48-0.52)	<.0001
≥30	0.25 (0.24-0.27)	<.0001	0.26 (0.24-0.28)	<.0001
Aboriginal				
No	1 (Ref)		1 (Ref)	
Yes	1.74 (1.68-1.79)	<.0001	1.56 (1.50-1.69)	<.0001
Year of test				
Per year later	0.97 (0.96-0.97)	<.0001	1.02 (1.02-1.02)	<.0001
Region				
Metro	1 (Ref)		1 (Ref)	
Rural	1.07 (1.02-1.12)	0.002	0.99 (0.95-1.04)	0.79
Remote	1.30 (1.25-1.34)	<.0001	1.00 (0.96-1.04)	0.87
Factors associated with gonorrhoea positivity				
	IRR	P-value	aIRR	P-value
Age group				
15-19	1.77 (1.65-1.90)	<.0001	1.42 (1.32-1.52)	<.0001
20-24	1 (Ref)		1 (Ref)	
25-29	0.70 (0.64-0.77)	<.0001	0.70 (0.64-0.76)	<.0001
≥30	0.44 (0.39-0.50)	<.0001	0.51 (0.45-0.58)	<.0001
Aboriginal				
No	1 (Ref)		1 (Ref)	
Yes	17.54 (16.17-19.01)	<.0001	11.80 (10.77-12.91)	<.0001
Year of test				
Per year later	0.87 (0.86-0.89)	<.0001	0.95 (0.94-0.96)	<.0001
Region				
Metro	1 (Ref)		1 (Ref)	
Rural	0.77 (0.63-0.93)	0.007	0.66 (0.55-0.78)	<.0001
Remote	5.65 (5.19-6.16)	<.0001	1.61 (1.49-1.74)	<.0001

300

Figure 1: Chlamydia positivity in women by year, age group and Aboriginality, 2001-2013





Figure 2: Gonorrhoea positivity in women by year, age group and Aboriginality: 2001-2013



304

305

306

- 308 1. Guy RJ, Kong F, Goller J, Franklin N, Bergeri I, Dimech W, et al. A new national
309 Chlamydia Sentinel Surveillance System in Australia: evaluation of the first stage of
310 implementation. *Communicable diseases intelligence quarterly report*. 2010;34(3):319-28.
- 311 2. Liu B, Donovan B, Parker J, Guy R, Hocking J, Kaldor JM, et al. Increasing chlamydia
312 diagnoses but little change in hospitalisations for ectopic pregnancy and infertility among
313 women in New South Wales from 2001 to 2008. *Sexual health*. 2012;9(4):355-9.
- 314 3. The Kirby Institute. HIV, viral hepatitis and sexually transmissible infections in Australia
315 Annual Surveillance Report 2015. The Kirby Institute, UNSW Australia, Sydney NSW 2052;
316 2015.
- 317 4. Graham S, Smith LW, Fairley CK, Hocking J. Prevalence of chlamydia, gonorrhoea,
318 syphilis and trichomonas in Aboriginal and Torres Strait Islander Australians: a systematic
319 review and meta-analysis. *Sexual health*. 2016;13(2):99-113.
- 320 5. Royal Australian College of General Practitioners. *Communicable diseases, STIs.*
321 *Guidelines for preventative activities in general practice*, 8th edition, East Melbourne
322 RACGP2012.
- 323 6. Government of Western Australia Department of Health. *Data Linkage Western*
324 *Australia 2016* [updated 26 February 2016. Available from: <http://www.datalinkage-wa.org.au/>.
- 325 7. Holman CD, Bass AJ, Rouse IL, Hobbs MS. Population-based linkage of health
326 records in Western Australia: development of a health services research linked database.
327 *Australian and New Zealand journal of public health*. 1999;23(5):453-9.
- 328 8. Christensen D, Davis G, Draper G, Mitrou F, McKeown S, Lawrence D, et al. Evidence
329 for the use of an algorithm in resolving inconsistent and missing Indigenous status in
330 administrative data collections. *Aust J Soc Issues*. 2014;49(4):423-43.
- 331 9. Government of Western Australia Department of Health. *Testing data for notifiable*
332 *sexually transmissible infections and blood-borne viruses in Western Australia, 2009 to 2013.*
333 Department of Health, Western Australia; 2014.
- 334 10. Government of Western Australia Department of Health. *Notification of infectious*
335 *diseases and related conditions 2016* [Available
336 from: [http://ww2.health.wa.gov.au/Articles/N_R/Notification-of-infectious-diseases-and-](http://ww2.health.wa.gov.au/Articles/N_R/Notification-of-infectious-diseases-and-related-conditions)
337 [related-conditions](http://ww2.health.wa.gov.au/Articles/N_R/Notification-of-infectious-diseases-and-related-conditions).
- 338 11. Regan DG, Wilson DP, Hocking JS. Coverage is the key for effective screening of
339 *Chlamydia trachomatis* in Australia. *The Journal of infectious diseases*. 2008;198(3):349-58.
- 340 12. Ali H, Guy RJ, Fairley CK, Wand H, Chen MY, Dickson B, et al. Understanding trends
341 in genital *Chlamydia trachomatis* can benefit from enhanced surveillance: findings from
342 Australia. *Sexually transmitted infections*. 2012;88(7):552-7.
- 343 13. O'Connor CC, Ali H, Guy RJ, Templeton DJ, Fairley CK, Chen MY, et al. High
344 chlamydia positivity rates in Indigenous people attending Australian sexual health services.
345 *The Medical journal of Australia*. 2014;200(10):595-8.
- 346 14. Graham S, Smith LW, Fairley CK, Hocking J. Prevalence of chlamydia, gonorrhoea,
347 syphilis and trichomonas in Aboriginal and Torres Strait Islander Australians: a systematic
348 review and meta-analysis. *Sexual health*. 2016().
- 349 15. Australian Government Department of Health. *Third National Sexually Transmissible*
350 *Infections Strategy 2014-2017*. 2014.
- 351 16. Donovan B, Dimech W, Ali H, Guy R, Hellard M. Increased testing for *Neisseria*
352 *gonorrhoeae* with duplex nucleic acid amplification tests in Australia: implications for
353 surveillance. *Sexual health*. 2015(12):48-50.
- 354 17. de Visser RO, Richters J, Rissel C, Badcock PB, Simpson JM, Smith AM, et al. Change
355 and stasis in sexual health and relationships: comparisons between the First and Second
356 Australian Studies of Health and Relationships. *Sexual health*. 2014;11(5):505-9.
- 357 18. Mercer CH, Tanton C, Prah P, Erens B, Sonnenberg P, Clifton S, et al. Changes in
358 sexual attitudes and lifestyles in Britain through the life course and over time: findings from

359 the National Surveys of Sexual Attitudes and Lifestyles (Natsal). Lancet.
360 2013;382(9907):1781-94.

361 19. Sonnenberg P, Clifton S, Beddows S, Field N, Soldan K, Tanton C, et al. Prevalence,
362 risk factors, and uptake of interventions for sexually transmitted infections in Britain: findings
363 from the National Surveys of Sexual Attitudes and Lifestyles (Natsal). Lancet.
364 2013;382(9907):1795-806.

365 20. O'Rourke KM, Fairley CK, Samaranayake A, Collignon P, Hocking JS. Trends in
366 Chlamydia positivity over time among women in Melbourne Australia, 2003 to 2007. Sexually
367 transmitted diseases. 2009;36(12):763-7.

368 21. Australian Bureau of Statistics. Western Australia Female Population Estimates, by
369 Health Region and Aboriginality (aged 15-34 years). 2008-2011.

370 22. Chow EPF, Fehler G, Read TRH, Tabrizi SN, Hocking JS, Denham I, et al. Gonorrhoea
371 notifications and nucleic acid amplification testing in a very low-prevalence Australian female
372 population. Med J Australia. 2015;202(6):321-3.

373 23. Government of Western Australia Department of Health. Guidelines for Managing
374 Sexually Transmitted Infections - WA 4th Edition 2015 [16/10/2015]. Available
375 from: [http://silverbook.health.wa.gov.au/Default.asp?PublicationID=1&ParentSectionID=P1&
376 SectionID=299](http://silverbook.health.wa.gov.au/Default.asp?PublicationID=1&ParentSectionID=P1&SectionID=299).

377


378

Appendix 1

To investigate if changes in the frequency of retesting could potentially bias the positivity trends, analyses were repeated with positivity calculated using only the first positive test from a woman in a particular year and excluding all subsequent tests in that year, and also from the only first test ever recorded for a woman

Figures A and B show the trends in chlamydia and gonorrhoea positivity when only the first test recorded each year was included and figures C and D when only the first ever test recorded was included. Among Aboriginal women a similar trend in chlamydia positivity was seen when only the first test recorded each year was included, although overall positivity was higher (Figure A) than in the main analysis (Figure 1). When analyses were restricted to only the first test ever recorded there was a small but significant increase in positivity among young Aboriginal women (IRR 1.02, 95%CI 1.01-1.04, $p=0.003$ and IRR 1.05, 1.02-1.07, $p=0.003$ among 15-19 and 20-24 year olds respectively) (Figure C). Among non-Aboriginal women the trends in chlamydia positivity were consistent when the different definitions of positivity were used suggesting that it is unlikely that the increase in positivity among young non-Aboriginal women is being driven by higher frequency of repeat testing in the later years. For gonorrhoea positivity, consistent trends in decreasing positivity were seen irrespective of the way positivity was calculated, in both Aboriginal and non-Aboriginal women (Figure B and D).


Figure A: Chlamydia positivity* in women by year, age group and Aboriginality, 2001-2013



*Positivity calculated from the first test recorded each year for each woman (a woman may contribute only one entry per year but multiple entries to the analysis)

380


Figure B: Gonorrhoea positivity* in women by year, age group and Aboriginality, 2001-2013



*Positivity calculated from the first test recorded each year for each woman (a woman may contribute only one entry per year but multiple entries to the analysis)


381

Figure C: Chlamydia positivity* in women by year, age group and Aboriginality, 2001-2013



*Positivity calculated from the first test recorded for each woman (each woman only contributes once to the analysis)

Figure D: Gonorrhoea positivity* in women by year, age group and Aboriginality, 2001-2013



*Positivity calculated from the first test recorded for each woman (each woman only contributes once to the analysis)