

2018

Effect of adjunctive ranitidine for antipsychotic-induced weight gain: A systematic review of randomized placebo-controlled trials

Xiaojing Gu

Rui Chen

Chen-Hui Sun

Wei Zheng

Xin-Hu Yang

See next page for additional authors

Follow this and additional works at: https://researchonline.nd.edu.au/med_article



Part of the [Medicine and Health Sciences Commons](#)

This article was originally published as:

Gu, X., Chen, R., Sun, C., Zheng, W., Yang, X., Wang, S., Ungvari, G. S., Ng, C., Golenkov, A., Lok, G., Li, L., Chow, I., Wang, F., & Xiang, Y. (2018). Effect of adjunctive ranitidine for antipsychotic-induced weight gain: A systematic review of randomized placebo-controlled trials. *Journal of International Medical Research*, 46 (1), 22-32.

<http://doi.org/10.1177/0300060517716783>

This article is posted on ResearchOnline@ND at https://researchonline.nd.edu.au/med_article/912. For more information, please contact researchonline@nd.edu.au.



Authors

Xiaojing Gu, Rui Chen, Chen-Hui Sun, Wei Zheng, Xin-Hu Yang, Shi-Bin Wang, Gabor S. Ungvari, Chee Ng, Andrei Golenkov, Grace Lok, Lu Li, Ines Chow, Fei Wang, and Yu-Tao Xiang

This is an Open Access article distributed in accordance with the Creative Commons Attribution-Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms provided the original work is properly cited and the use is non-commercial.

See: <https://creativecommons.org/licenses/by-nc/4.0/>

This article originally published in *Journal of International Medical Research* available at:

<http://dx.doi.org/10.1177/0300060517716783>

No changes have been made to this article.

Gu. X., Chen, R., Sun, C., Zheng, W., Yang, X., Wang, G. ... Xiang, Y. (2018). Effect of adjunctive ranitidine for antipsychotic-induced weight gain: A systematic review of randomized placebo-controlled trials. *Journal of International Medical Research*, 46(1), 22-32. doi : 10.1177/0300060517716783

Effect of adjunctive ranitidine for antipsychotic-induced weight gain: A systematic review of randomized placebo-controlled trials

Xiao-Jing Gu^{1,*}, Rui Chen^{2,*}, Chen-Hui Sun^{3,*},
Wei Zheng^{4,*}, Xin-Hu Yang⁴, Shi-Bin Wang⁵,
Gabor S. Ungvari^{6,7}, Chee H. Ng⁸,
Andrei Golenkov⁹, Grace K.I. Lok⁵, Lu Li⁵,
Ines H.I. Chow⁵, Fei Wang⁵ and Yu-Tao Xiang⁵

Abstract

This study was a meta-analysis of randomized controlled trials (RCTs) of ranitidine as an adjunct for antipsychotic-induced weight gain in patients with schizophrenia. RCTs reporting weight gain or metabolic side effects in patients with schizophrenia were included. Case reports/series, non-randomized or observational studies, reviews, and meta-analyses were excluded. The primary outcome measures were body mass index (BMI) (kg/m^2) and body weight (kg). Four RCTs with five study arms were identified and analyzed. Compared with the control group, adjunctive ranitidine was associated with marginally significant reductions in BMI and body weight. After removing an outlier study for BMI, the effect of ranitidine remained significant. Adjunctive ranitidine outperformed the placebo in the negative symptom score of the Positive and Negative Syndrome Scale. Although ranitidine was associated with less frequent drowsiness, other adverse events were similar between the two groups. Adjunctive ranitidine appears to be an effective and

¹The Second Affiliated Hospital of Xinxiang Medical University Henan Mental Hospital, Henan, China

²Beijing Key Laboratory of Mental Disorders, Beijing Anding Hospital, Capital Medical University, Beijing, China

³Qingdao Mental Health Center, Shandong, China

⁴The Affiliated Brain Hospital of Guangzhou Medical University (Guangzhou Huiai Hospital), Guangzhou, China

⁵Unit of Psychiatry, Faculty of Health Sciences, University of Macau, Macao SAR, China

⁶The University of Notre Dame Australia/Marian Centre, Perth, Australia

⁷School of Psychiatry & Clinical Neurosciences, University of Western Australia, Perth, Australia

⁸Department of Psychiatry, University of Melbourne, Melbourne, Victoria, Australia

⁹Department of Psychiatry and Medical Psychology, Chuvash State University, Cheboksary, Russia

*These authors contributed equally to this work.

Corresponding author:

Yu-Tao Xiang, 3/F, Building E12, Faculty of Health Sciences, University of Macau, Avenida da Universidade, Taipa, Macau SAR 999078, China.

Email: xyutly@gmail.com



safe option for reducing antipsychotic-induced weight gain and improving negative symptoms in patients with schizophrenia. Larger RCTs are warranted to confirm these findings.

Trial registration: PROSPERO: CRD42016039735

Keywords

Schizophrenia, ranitidine, weight gain, negative symptom, meta-analysis, antipsychotics

Date received: 20 January 2017; accepted: 1 June 2017

Introduction

Antipsychotic (AP)-induced weight gain is common in patients with schizophrenia and has received increased attention in recent decades.^{1–7} In one study, up to 75% of patients with a first episode of schizophrenia receiving APs experienced weight gain of $\geq 7\%$.⁸ AP-induced weight gain not only increases the risk of diabetes mellitus, cardiovascular disease, and mortality but also contributes to poor treatment adherence and low quality of life.^{2,3,9–11}

Histamine H2 receptors play an important role in the mechanism of AP-induced weight gain, together with the serotonergic, noradrenergic, and histaminergic systems.¹² Ranitidine, a non-imidazole H2 blocker, is an inexpensive medication with a low risk of adverse effects because of its negligible effects on muscarinic, nicotinic, adrenergic, and H1 receptors.¹³ Although several randomized controlled trials (RCTs)^{13–16} have examined the efficacy and safety of adjunctive ranitidine for AP-induced weight gain in patients with schizophrenia, the results were mixed.

To the best of our knowledge, no systematic reviews or meta-analyses on ranitidine as an adjunctive treatment for AP-induced weight gain in patients with schizophrenia have been published. Therefore, we performed the present meta-analysis of RCTs of adjunctive ranitidine for all AP-induced weight gain in patients with schizophrenia. We also included recent RCTs published in Chinese-language journals

that may not be widely known on an international basis.

Methods

Search strategy and selection criteria

Two reviewers independently and systematically searched the PubMed, PsycINFO, Embase, Cochrane Library, Chinese Journal Net, WanFang, and China Biology Medicine databases for articles regarding adjunctive ranitidine for schizophrenia from inception of each database until 10 October 2016. The keywords used for the search were (zantic OR ranitidine OR zantic OR Zantac) AND (schizophrenic disorder OR disorder, schizophrenic OR schizophrenic disorders OR schizophrenia OR dementia praecox). We also hand-searched reference lists from relevant review articles for additional studies and contacted the authors for more information if necessary.

The following criteria were used according to the *PICOS* acronym. Participants (*P*): adult patients (≥ 18 years of age) with schizophrenia using any diagnostic criteria. Intervention (*I*): ranitidine plus APs. Comparison (*C*): APs plus placebo or AP monotherapy. Outcomes (*O*): efficacy and safety. Study design (*S*): RCT reporting body weight or metabolic adversities as primary or secondary outcomes. Case reports/series, non-randomized or observational studies, reviews, and meta-analyses were excluded.

Outcome measures

The primary outcome measures were body mass index (BMI) (kg/m^2) and body weight (kg). Key secondary outcomes were clinical improvement assessed by the Positive and Negative Syndrome Scale (PANSS)¹⁷ or the Brief Psychiatric Rating Scale,¹⁸ discontinuation rate, and adverse drug reactions (ADRs).

Data extraction

Data were identified, checked, extracted, and analyzed by two independent reviewers. Furthermore, outcomes based on intention-to-treat analysis, if available, were recorded. Any inconsistencies were discussed with and resolved by a third reviewer.

Statistical methods

According to the guidelines of the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement,¹⁹ Review Manager (RevMan) Version 5.3 (<http://tech.cochrane.org/revman/>) was used to perform the meta-analysis. For meta-analytic pooling of continuous and dichotomous outcomes, the inverse variance method and Mantel–Haenszel test were used to present weighted mean differences (WMDs) and risk ratios (RRs) with their 95% confidence intervals (CIs), respectively. When the RR was significant, the number needed to treat or number needed to harm (NNH) was calculated by dividing 1 by the risk difference. Each missing standard deviation (SD) was replaced by the average SD from other RCTs that used the same medication.²⁰ To compensate for study heterogeneity, a random-effects model was used in all meta-analyzable data.²¹ Both I^2 and chi-square statistics were used to identify heterogeneity. When heterogeneity (chi-squared $P < 0.1$ and $I^2 > 50\%$) was present for BMI, a sensitivity analysis was conducted to examine the credibility of the BMI

change by excluding one study¹⁶ with an outlying effect size (ES) of less than -1.5 (i.e. more than 1.5 SD superiority of ranitidine). In addition, one subgroup analysis was performed (Chinese versus non-Chinese patients). One study¹³ with three treatment arms compared the combination of ranitidine and two different doses of APs with a control group. Half of the patients were assigned to each ranitidine arm to avoid inflating the number of patients in the control group. Publication bias was assessed using funnel plots and Egger's test.²² All analyses were two-tailed, with alpha set at 0.05.

Assessment of reporting biases

According to the recommendations of the Cochrane Collaboration, the Cochrane risk of bias was used to assess the methodological quality of RCTs (Supplementary Figure 1). Their domain was rated as “high risk,” “unclear risk,” or “low risk.”²³

The Grading of Recommendations Assessment, Development and Evaluation (GRADE) system was used to rate each study as “very low,” “low,” “moderate,” or “high” with respect to the quality of evidence of adjunctive ranitidine versus placebo. Furthermore, the Jadad scale (Table 1) ranging from 0 to 5 was used to judge the quality of each RCT.²⁴

Results

Results of the search

Figure 1 presents a flow chart of article selection from the English ($n = 114$) and Chinese databases ($n = 11$). The full text of one RCT²⁵ published in Spanish could not be obtained. In total, four RCTs^{13–16} were eligible and included in this meta-analysis.

Study characteristics

The four RCTs^{13–16} with five study arms ($n = 315$) comprised three double-blinded

Table 1. Study, patient, and treatment characteristics.

Study	Number of patients	Blinding	Trial duration, weeks	Setting, %	Diagnosis, %	Diagnostic criteria	Illness duration, years	Age at baseline, years (range)	Male sex, %	Control group:		Jadad score	
										Dosage in mg/d, mean (range)	Intervention group: Dosage in mg/d, mean (range)		
Liang 2016 (China)	T: 120	DB	ITT	Inpatients (100)	Sz (100)	CCMD-3	<5.0	33.0 (18–45)	52.5	OLA:	OLA:	4	
	M: 60									NR (5–30)	NR (5–30)		150 (FD)
	A: 60												
Mehta and Ram 2014 (India)	T: 75	OL	ITT	Inpatients (100)	Sz (100)	ICD-10	5.0 (18–60)	31.2	89.3	OLA:	OLA:	3	
	M: 25									25.7 (10–30)	24.6 (10–30)		150 (FD)
	A: 50												
Ranjbar et al. 2013 (Iran)	T: 52	DB	ITT	Inpatients (100)	Sz (NR)	DSM-IV	NR	38.1 (NR)	63.5	OLA:	OLA:	5	
	M: 27									NR (NR)	NR (NR)		600 (FD)
	A: 25												
Sun et al. 2007 (China)	T: 68	DB	OC	Inpatients (100)	Sz (100)	CCMD-3	1.6 (19–48)	28.6	56.9	OLA:	OLA:	4	
	M: 33									NR (10–20)	NR (10–20)		300 (FD)
	A: 35												

A = augmentation; CCMD-3 = China's Mental Disorder Classification and Diagnosis Standard, 3rd edition; DSM-IV = Diagnostic and Statistical Manual of Mental Disorders, 4th edition; DB = double blind; FD = fixed dosage; ICD-10 = International Statistical Classification of Diseases and Related Health Problems, 10th revision; ITT = intention to treat; M = monotherapy; NR = not reported; OL = open label; OC = observed cases; OLA = olanzapine; RAN = ranitidine; Sz = schizophrenia; SZA = schizoaffective disorder; SZD = schizophreniform disorder; T = total.




Figure 1. PRISMA flow diagram. RCT, randomized controlled trial.

trials ($n=240$) and one open label study ($n=75$) comparing an adjunctive ranitidine group ($n=170$) and a control group ($n=145$). The weighted mean treatment duration was 15.8 weeks (range = 8–24 weeks) (Table 1). Two RCTs were conducted in China ($n=188$), one was conducted in India ($n=75$), and one was conducted in Iran ($n=52$).

Patient characteristics

The weighted mean age was 32.4 years (range = 28.6–38.1 years), the mean percentage

of male patients was 64% (range = 52.5%–89.3%), and the weighted mean illness duration (according to the available data in two RCTs^{13,15}) was 3.2 years (range = 1.6–5.0 years) (Table 1). All RCTs were conducted in inpatient settings.

Treatment characteristics

The weighted mean dosage of ranitidine was 339.5 mg/day (range = 150–600 mg/day). All included studies involved only patients who received olanzapine. The weighted mean dosage of olanzapine was 19.1 mg/day

Table 2. GRADE analyses: Adjunctive ranitidine for antipsychotic-induced weight gain.

Design	N (arms)	Risk of bias	Inconsistency	Indirectness	Imprecision	Publication bias	Large effect	Overall quality of evidence ^a
Body weight (kg)	312 (5)	None	Serious ^b	None	None	Serious ^c	None	++/-/-/-; Low
BMI (kg/m ²)	260 (4)	None	Serious ^b	None	None	Serious ^c	None	++/-/-/-; Low
PANSS total score	75 (2)	Serious ^d	None	None	None	Serious ^c	None	++/-/-/-; Low
PANSS positive symptom score	75 (2)	Serious ^d	None	None	None	Serious ^c	None	++/-/-/-; Low
PANSS negative symptom score	75 (2)	Serious ^d	None	None	None	Serious ^c	None	++/-/-/-; Low
PANSS general symptom score	75 (2)	Serious ^d	None	None	None	Serious ^c	None	++/-/-/-; Low
Drowsiness	165 (3)	None	None	None	None	Serious ^e	None	++/+/+/-; Moderate

BMI = body mass index; GRADE = Grading of Recommendations Assessment, Development and Evaluation; PANSS = Positive and Negative Syndrome Scale.

^aGRADE Working Group grades of evidence: High quality = further research is very unlikely to change our confidence in the estimate of effect. Moderate quality = further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate. Low quality = further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate. Very low quality = we are very uncertain about the estimate.

^bAll studies that reported having a serious inconsistency had $I^2 > 50\%$.

^cFor continuous outcomes, $N < 400$.

^dAll studies that reported having a serious bias used an open label method and only mentioned random allocation without describing the method.

^eFor dichotomous outcomes, $N < 300$.

(range = 5–30 mg/day) according to the available data in three RCTs.^{13,15,16}

Quality assessment

All four RCTs mentioned “randomized allocation” with a specific description, while the allocation concealment method was rated as “unclear” in one RCT and “high risk” in another RCT. Three RCTs were double-blinded trials, and one RCT was an open label study. Regarding incomplete outcome data, one RCT reported loss to follow-up but failed to use intention-to-treat analysis. Furthermore, two RCTs employed a protocol registration and were rated as “low risk” regarding selective reporting (Supplementary Figure 1). The quality of evidence presented for the primary and secondary outcomes using the GRADE approach ranged from “low” (86%) to “moderate” (14%) (Table 2).

The weighted mean total Jadad score of the four RCTs was 3.9 (range = 3–5), and all studies were rated as high-quality (Jaded score of ≥ 3) (Table 1).

Primary outcomes

Compared with the control group, ranitidine was associated with a marginally significant decrease in the BMI (4 RCTs with 5 study arms, $n = 312$; WMD: -1.08 kg/m²; 95%CI: $-2.15, -0.01$; $P = 0.05$; $I^2 = 94\%$) (Figure 2). However, the BMI change was significant even after removing an outlier study¹⁶ ($n = 120$; $ES < -1.5$; WMD: -0.58 kg/m²; 95%CI: $-1.08, -0.07$; $P = 0.02$; $I^2 = 64\%$). Moreover, the subgroup analysis revealed that statistical significance was lost for studies of both non-Chinese patients (2 RCTs with 3 study arms, $n = 127$) and Chinese patients (2 RCTs, $n = 185$). Furthermore, patients taking




Figure 2. Ranitidine for antipsychotic-induced weight gain: Forest plot for changes in body weight and body mass index.

BMI, body mass index; SD, standard deviation; IV, interval variance; CI, confidence interval.

ranitidine did not lose weight to a statistically significant degree but only showed a trend (3 RCTs with 4 study arms, $n = 260$; WMD: -1.54 kg; 95%CI: $-3.13, 0.04$; $I^2 = 78\%$) (Figure 2) compared with the control group. Because only four RCTs with five study arms were included for assessment of the primary outcomes, the presence of publication bias regarding weight change and BMI could not be determined by performing a funnel plot or Egger's test (< 10 trials).²⁶

Secondary outcomes

With respect to clinical outcomes, only one RCT with two study arms used the PANSS. The adjunctive ranitidine group outperformed the control group in terms of PANSS negative symptom scores (1 RCT, $n = 75$; WMD: -1.95 ; 95%CI: $-3.62, -0.28$; $P = 0.02$; $I^2 = 0\%$) (Figure 3), but not in terms of PANSS total, positive, or general symptom scores (1 RCT, $n = 75$; WMD: -0.44 to 0.77 ; 95%CI: $-3.88, 2.99$; $I^2 = 0\%$) (Figure 3).

In terms of ADRs, adjunctive ranitidine was associated with less frequent drowsiness than in the control group (3 RCTs, $n = 165$;

RR: 0.55; 95%CI: 0.35, 0.86; $P = 0.008$; $I^2 = 0\%$; NNH = 6; 95%CI = 3, 100) (Figure 4). Meta-analyses of akathisia, rigidity, tremor, dry mouth, headache, and constipation showed no significant group differences (Figure 4).

One RCT¹⁵ reported a discontinuation rate of 6% (2/35) in the ranitidine group versus 3% (1/33) in the control group. The remaining RCTs did not report the discontinuation rate.

Discussion

To the best of our knowledge, this is the first meta-analysis to examine the effect of adjunctive ranitidine on AP-induced weight gain. The adjunctive ranitidine group outperformed the control group in terms of reductions in weight gain and improvements in negative symptoms. Furthermore, ranitidine was well tolerated. These positive effects of ranitidine support the hypothesis that the histamine H2 receptor is a possible mediator of eating behavior and weight regulation.²⁷ Importantly, the significance the effect remained when the one outlier ($ES < -1.5$)¹⁶ was excluded from the analysis. The lack of




Figure 3. Ranitidine for antipsychotic-induced weight gain: Forest plot for clinical efficacy assessed by changes in the Positive and Negative Syndrome Scale (PANSS) score. SD, standard deviation; IV, interval variance; CI, confidence interval.

significant results regarding Chinese and non-Chinese participants in the subgroup analysis could have been due to the small sample size, which reduced the power to detect statistically significant results.

In this study, the ranitidine group was superior to the control group with respect to improvements in negative symptoms (WMD = -1.95). This may have been due to the direct drug receptor activity of ranitidine itself.¹³ Ranitidine has been shown to be relatively safe and well tolerated in patients with schizophrenia. Drowsiness (NNH = 6) was less frequent in the ranitidine group than in the control group, and no group difference in other ADRs was evident. Topiramate and metformin may also improve AP-related metabolic ADRs.⁶ However, we could not locate any head-to-head trials comparing adjunctive ranitidine with topiramate/metformin in patients with schizophrenia.

Limitations

This study had several limitations. First, there was significant heterogeneity of the results in terms of the primary outcomes. However, a random-effects model was employed to provide a conservative estimate for all meta-analyzable outcomes. Second, only a few studies were available for the meta-analysis because the use of ranitidine for weight loss is off-label. The inclusion of only 4 RCTs involving 315 patients and the limited or incomplete information lessen the confidence of the results and limit more comprehensive data exploration, such as meta-regression analyses. Third, although all four RCTs were rated as high-quality using the Jadad scale,²⁴ 86% were rated as “low” using the GRADE approach, especially for BMI and weight change. Furthermore, publication bias for primary outcomes could not be examined because of




Figure 4. Ranitidine for antipsychotic-induced weight gain: Forest plot for adverse drug reactions. MH, Mantel–Haenszel.

the limited number of studies.²⁸ Fourth, because the ranitidine dosages varied from 150 to 600 mg/day, the dose–response effect of ranitidine in weight loss could not be further evaluated. Fifth, all RCTs used

olanzapine as the baseline AP; therefore, the results could not be generalized to other APs. Furthermore, the >24-week long-term effects of adjunctive ranitidine on body weight could not be investigated. Finally,

other metabolic indices including the lipid profile, insulin resistance, and leptin concentration were not recorded in the included studies.

Conclusions

Adjunctive ranitidine appears to be an effective and safe option for treating weight gain and negative symptoms in patients with schizophrenia. Because of the small number of studies available for this meta-analysis, the results should be regarded as preliminary. The long-term effects of ranitidine on weight change and negative symptoms need to be examined in further studies with improved methodology. In addition, the effect of ranitidine on weight gain in patients with other psychiatric disorders should be examined in systematic reviews.

Acknowledgements

The study was supported by the University of Macau (SRG2014-00019-FHS; MYRG2015-00230-FHS; MYRG2016-00005-FHS) and Affiliated Brain Hospital of Guangzhou Medical University (22016YFC0906302; 2014Y2-00105; 81671334). The University of Macau and Affiliated Brain Hospital of Guangzhou Medical University had no role in the study design, generation or interpretation of the results, or publication of the study. We wish to thank Prof. Phil Wiffen (Training Director of UK Cochrane Center) and Dr. Jun Xia (Cochrane Schizophrenia Group) for their systematic review courses.

Declaration of conflicting interests

The authors declare that there is no conflict of interest.

Funding

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

References

1. Correll CU, Sikich L, Reeves G, et al. Metformin for antipsychotic-related weight gain and metabolic abnormalities: when, for whom, and for how long? *Am J Psychiatry* 2013; 170: 947–952.
2. Maayan L, Vakhrusheva J and Correll CU. Effectiveness of medications used to attenuate antipsychotic-related weight gain and metabolic abnormalities: a systematic review and meta-analysis. *Neuropsychopharmacology* 2010; 35: 1520–1530.
3. Zheng W, Li XB, Tang YL, et al. Metformin for weight Gain and metabolic abnormalities associated with antipsychotic treatment: meta-analysis of randomized placebo-controlled trials. *J Clin Psychopharmacol* 2015; 35: 499–509.
4. Wu RR, Zhang FY, Gao KM, et al. Metformin treatment of antipsychotic-induced dyslipidemia: an analysis of two randomized, placebo-controlled trials. *Mol Psychiatry* 2016; 21: 1537–1544.
5. Liu Z, Zheng W, Gao S, et al. Metformin for treatment of clozapine-induced weight gain in adult patients with schizophrenia: a meta-analysis. *Shanghai Arch Psychiatry* 2015; 27: 331–340.
6. Zheng W, Xiang YT, Xiang YQ, et al. Efficacy and safety of adjunctive topiramate for schizophrenia: a meta-analysis of randomized controlled trials. *Acta Psychiatr Scand* 2016; 134: 385–398.
7. Zheng W, Zheng YJ, Li XB, et al. Efficacy and safety of adjunctive aripiprazole in schizophrenia: meta-analysis of randomized controlled trials. *J Clin Psychopharmacol* 2016; 36: 628–636.
8. Curtis J, Watkins A, Rosenbaum S, et al. Evaluating an individualized lifestyle and life skills intervention to prevent antipsychotic-induced weight gain in first-episode psychosis. *Early Interv Psychiatry* 2016; 10: 267–276.
9. Mizuno Y, Suzuki T, Nakagawa A, et al. Pharmacological strategies to counteract antipsychotic-induced weight gain and metabolic adverse effects in schizophrenia: a systematic review and meta-analysis. *Schizophr Bull* 2014; 40: 1385–1403.

10. Poyurovsky M, Isaacs I, Fuchs C, et al. Attenuation of olanzapine-induced weight gain with reboxetine in patients with schizophrenia: a double-blind, placebo-controlled study. *Am J Psychiatry* 2003; 160: 297–302.
11. Choi YJ. Efficacy of adjunctive treatments added to olanzapine or clozapine for weight control in patients with schizophrenia: a systematic review and meta-analysis. *ScientificWorldJournal* 2015; 2015: 970730.
12. Baptista T, Kin NM, Beaulieu S, et al. Obesity and related metabolic abnormalities during antipsychotic drug administration: mechanisms, management and research perspectives. *Pharmacopsychiatry* 2002; 35: 205–219.
13. Mehta VS and Ram D. Efficacy of ranitidine in olanzapine-induced weight gain: a dose-response study. *Early Interv Psychiatry* 2014; 10: 522–527.
14. Ranjbar F, Ghanepour A, Sadeghi-Bazargani H, et al. The effect of ranitidine on olanzapine-induced weight gain. *Biomed Res Int* 2013; 2013: 639391.
15. Sun J, Xi XD, Li Y, et al. The effect of ranitidine on olanzapine-induced weight gain and metabolic dysfunction [In Chinese]. *Chinese Journal of Nervous and Mental Diseases* 2007; 33: 560–562.
16. Liang MJ. Effects of ranitidine on the glucose and lipid metabolism level and body mass of patients with olanzapine in treatment of schizophrenia [In Chinese]. *Evaluation and Analysis of Drug-use in Hospitals of China* 2016; 16: 433–436.
17. Kay SR, Fiszbein A and Opler LA. The positive and negative syndrome scale (PANSS) for schizophrenia. *Schizophr Bull* 1987; 13: 261–276.
18. Overall JE and Gorham DR. The brief psychiatric rating-scale. *Psychol Rep* 1962; 10: 799–812.
19. Moher D, Liberati A, Tetzlaff J, et al. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Ann Intern Med* 2009; 151: 264–269, W264.
20. Leucht S, Komossa K, Rummel-Kluge C, et al. A meta-analysis of head-to-head comparisons of second-generation antipsychotics in the treatment of schizophrenia. *Am J Psychiatry* 2009; 166: 152–163.
21. DerSimonian R and Laird N. Meta-analysis in clinical trials. *Control Clin Trials* 1986; 7: 177–188.
22. Egger M, Davey Smith G, Schneider M, et al. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997; 315: 629–634.
23. Higgins J and Higgins J. *Cochrane Handbook for Systematic Reviews of Interventions*. Chichester, UK: John Wiley & Sons, 2008.
24. Jadad AR, Moore RA, Carroll D, et al. Assessing the quality of reports of randomized clinical trials: is blinding necessary? *Control Clin Trials* 1996; 17: 1–12.
25. Lopez-Mato A, Rovner J, Illa G, et al. [Randomized, open label study on the use of ranitidine at different doses for the management of weight gain associated with olanzapine administration]. *Vertex* 2003; 14: 85–96. [in Spanish, English Abstract].
26. Sterne JA, Sutton AJ, Ioannidis JP, et al. Recommendations for examining and interpreting funnel plot asymmetry in meta-analyses of randomised controlled trials. *BMJ* 2011; 343: d4002.
27. Doi T, Sakata T, Yoshimatsu H, et al. Hypothalamic neuronal histamine regulates feeding circadian rhythm in rats. *Brain Res* 1994; 641: 311–318.
28. Reynolds GP and Kirk SL. Metabolic side effects of antipsychotic drug treatment—pharmacological mechanisms. *Pharmacol Ther* 2010; 125: 169–179.