

2018

Cerebral small vessel disease burden is associated with poststroke depressive symptoms: A 15-month prospective study

Yan Liang

Yang-Kun Chen

Vincent C. Mok

De-Feng Wang

Gabor S. Ungvari

The University of Notre Dame Australia, Gabor.Ungvari@nd.edu.au

See next page for additional authors

Follow this and additional works at: https://researchonline.nd.edu.au/med_article



Part of the [Medicine and Health Sciences Commons](#)

This article was originally published as:

Liang, Y., Chen, Y., Mok, V. C., Wang, D., Ungvari, G. S., Chu, W. C., Kang, H., & Tang, W. (2018). Cerebral small vessel disease burden is associated with poststroke depressive symptoms: A 15-month prospective study. *Frontiers in Aging Neuroscience*, 10.

Original article available here:

<https://doi.org/10.3389/fnagi.2018.00046>

This article is posted on ResearchOnline@ND at
https://researchonline.nd.edu.au/med_article/916. For more
information, please contact researchonline@nd.edu.au.



Authors

Yan Liang, Yang-Kun Chen, Vincent C. Mok, De-Feng Wang, Gabor S. Ungvari, Winnie C. Chu, Hee-Ju Kang, and Wai-Kwong Tang

This is an Open Access article distributed in accordance with the Creative Commons Attribution 4.0 International (CC BY 4.0) License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

See: <https://creativecommons.org/licenses/by/4.0/>

This article originally published in *Frontiers in Aging Neuroscience* available at:

<https://doi.org/10.3389/fnagi.2018.00046>

Liang, Y., Chen, Y., Mok, V.C., Wang, D., Ungvari, G.S., Chu, W.C., Kang, H., and Tang, W. (2018). Cerebral small vessel disease burden is associated with poststroke depressive symptoms: A 15-month prospective study. *Frontiers in Aging Neuroscience*, 10. doi: 10.3389/fnagi.2018.00046



Cerebral Small Vessel Disease Burden Is Associated With Poststroke Depressive Symptoms: A 15-Month Prospective Study

Yan Liang¹, Yang-Kun Chen², Vincent Chung-Tong Mok³, De-Feng Wang⁴, Gabor S. Ungvari⁵, Winnie Chiu-Wing Chu⁴, Hee-Ju Kang⁶ and Wai-Kwong Tang^{1,7*}

¹ Department of Psychiatry, The Chinese University of Hong Kong, Hong Kong, Hong Kong, ² Department of Neurology, Dongguan People's Hospital, Dongguan, China, ³ Department of Medicine and Therapeutics, The Chinese University of Hong Kong, Hong Kong, Hong Kong, ⁴ Department of Imaging and Interventional Radiology, The Chinese University of Hong Kong, Hong Kong, Hong Kong, ⁵ Graylands Hospital, The University of Notre Dame Australia, Perth, WA, Australia, ⁶ Department of Psychiatry, Chonnam National University Hospital, Gwangju, South Korea, ⁷ Shenzhen Research Institute, The Chinese University of Hong Kong, Shenzhen, China

OPEN ACCESS

Edited by:

Aurel Popa-Wagner,
University of Rostock, Germany

Reviewed by:

Raluca Sandu Vintilescu,
University of Medicine and Pharmacy
of Craiova, Romania
Danny J. J. Wang,
University of Southern California,
United States

*Correspondence:

Wai-Kwong Tang
tangwk@cuhk.edu.hk

Received: 28 November 2017

Accepted: 12 February 2018

Published: 28 February 2018

Citation:

Liang Y, Chen Y-K, Mok VC-T, Wang D-F, Ungvari GS, Chu WC-W, Kang H-J and Tang W-K (2018) Cerebral Small Vessel Disease Burden Is Associated With Poststroke Depressive Symptoms: A 15-Month Prospective Study. *Front. Aging Neurosci.* 10:46. doi: 10.3389/fnagi.2018.00046

Objective: All types of cerebral small vessel disease (SVD) markers including lacune, white matter hyperintensities (WMH), cerebral microbleeds, and perivascular spaces were found to be associated with poststroke depressive symptoms (PDS). This study explored whether the combination of the four markers constituting an overall SVD burden was associated with PDS.

Methods: A cohort of 563 patients with acute ischemic stroke were followed over a 15-month period after the index stroke. A score of ≥ 7 on the 15-item Geriatric Depression Scale was defined as clinically significant PDS. Scores of the four SVD markers ascertained on magnetic resonance imaging were summed up to represent total SVD burden. The association between SVD burden and PDS was assessed with generalized estimating equation models.

Results: The study sample had a mean age of 67.0 ± 10.2 years and mild-moderate stroke [National Institutes of Health Stroke Scale score: 3, interquartile, 1–5]. PDS were found in 18.3%, 11.6%, and 12.3% of the sample at 3, 9, and 15 months after stroke, respectively. After adjusting for demographic characteristics, vascular risk factors, social support, stroke severity, physical and cognitive functions, and size and locations of stroke, the SVD burden was associated with an increased risk of PDS [odds ratio = 1.30; 95% confidence interval = 1.07–1.58; $p = 0.010$]. Other significant predictors of PDS were time of assessment, female sex, smoking, number of acute infarcts, functional independence, and social support.

Conclusion: SVD burden was associated with PDS examined over a 15-month follow-up in patients with mild to moderate acute ischemic stroke.

Keywords: depressive symptoms, stroke, poststroke depression, cerebral small vessel disease, white matter hyperintensities, lacune, cerebral microbleeds, enlarged perivascular spaces

INTRODUCTION

Poststroke depression (PSD) is associated with higher risk of mortality, disability, and poor quality of life (Towfighi et al., 2017). About one in three stroke survivors present with PSD at any time after stroke (Towfighi et al., 2017). Understanding the underlying pathophysiology of PSD may facilitate developing better management strategies. Findings concerning the relationship between stroke locations and PSD have been inconsistent (Carson et al., 2000; Wei et al., 2015), which challenges the hypothesis that PSD is caused by a single lesion in the brain due to stroke (Stern and Bachman, 1991).

Cerebral small vessel disease (SVD) is a composite term for a variety of chronic cerebral microvascular lesions resulting from aging, vascular risk factors or unknown etiologies (Wardlaw et al., 2013). SVD is common in aging and stroke populations (Wardlaw et al., 2013). The most characteristic markers of SVD on magnetic resonance imaging (MRI) are WMH, lacune, cerebral microbleeds, and enlarged perivascular spaces (Wardlaw et al., 2013). A “SVD score” that includes the above four MRI markers was recently proposed to quantify the global damage to the small vessels of the brain caused by SVD (Staals et al., 2014). The SVD burden is associated with cognitive decline in hypertension (Uiterwijk et al., 2016), poststroke disability (Arba et al., 2017a), recurrence of stroke (Lau et al., 2017), poor quality of life of stroke survivors (Liang et al., 2017), and depression after lacunar stroke (Zhang X. et al., 2017).


Individual SVD markers have been shown to be associated with PSD (Santos et al., 2009a; Tang et al., 2014). Only one study has reported an association between SVD burden and poststroke depressive symptoms (PDS) 3 months after lacunar stroke (Zhang X. et al., 2017). However, the impact of SVD burden on the long-term prevalence of PSD has never been investigated. Similarly, the role of SVD burden in the development of PSD in a cohort comprising all subtypes of acute ischemic stroke is unclear. Hence, this study examined the association between SVD burden and poststroke depressive symptoms (PDS) during the first 15 months following an acute ischemic stroke.

MATERIALS AND METHODS

Study Population

The recruitment of the study sample is depicted in **Figure 1** where T0–T3 refer to assessment points at baseline, 3, 9, and 15 months after the index stroke, respectively. Patients with first-ever or recurrent acute ischemic stroke were consecutively screened following their admission to the Stroke Unit, Prince of

Abbreviations: BI, Barthel Index; CI, confidence interval; GDS, Geriatric Depression Scale; GEE, generalized estimating equation; IQR, interquartile ranges; LSNS, Lubben Social Network Scale; MMSE, Mini-Mental State Examination; MRI, Magnetic Resonance Imaging; NIHSS, National Institutes of Health Stroke Scale; OR, odds ratio; PDS, poststroke depressive symptoms; PSD, poststroke depression; SD, standard deviations; SVD, small vessel disease; WMH, white matter hyperintensities.



Wales Hospital, Hong Kong, between January 2010 and October 2015. Due to the limited access to MRI, only 2,048 (32.1%) of all admitted stroke patients underwent an MRI examination within 1 week of the onset of stroke. A further 1,339 (62.0%) were excluded because they (1) refused to participate or were lost to follow-up; (2) had a history of psychiatric or neurologic disorders; (3) had severe comorbid medical conditions, such as multi-functional organs failure, cancer or severe systemic infection; (4) had severe aphasia, auditory or visual impairments; (5) deceased before the 15-month assessment; (6) had a recurrent stroke during the 15-month follow-up; and (7) were not of Chinese ethnicity.

The study protocol was approved by the Joint Chinese University of Hong Kong New Territories East Cluster Clinical Research Ethics Committee. All participants gave written informed consent.

Demographic and Clinical Data

Within 1 week of the index admission (T0), demographic (age, sex, and year of education) and clinical data on vascular risk factors (smoking, history of hypertension, diabetes mellitus, hyperlipidaemia, stroke, atrial fibrillation, and ischemic heart disease) were collected. Baseline stroke severity was assessed with the National Institutes of Health Stroke Scale (NIHSS) (Brott et al., 1989) by a nurse from the Stroke Unit. Physical and cognitive functions and level of social

support were assessed by a research assistant with the Barthel Index (BI) (Barthel, 1965), the Mini-Mental State Examination (MMSE) (Chiu et al., 1994), and the Lubben Social Network Scale (LSNS) (Lubben, 1988), respectively at T1, T2, and T3.

Study Outcomes

The primary study outcome was PDS at either T1, T2, or T3. All patients were screened for PDS with the 15-item Geriatric Depression Scale (GDS) by a psychiatrist (W-KT). Clinically significant PDS suspect for an ongoing depressive illness was defined by a score 7 or above on the GDS (Tang et al., 2004a). The 15-item GDS has good sensitivity (89%) and specificity (73%) in screening for PDS in a Hong Kong stroke population (Tang et al., 2004a).

Neuroimaging

All patients were scanned on a 1.5-T MRI system (Sonata; Siemens Medical Systems Inc., Germany) with diffusion weighted, gradient echo, T1- and T2-weighted and fluid attenuated inversion recovery sequences. The MRI parameters were described in detail elsewhere (Liang et al., 2017). A qualified neurologist (YL) blind to all demographic and clinical information reviewed the MRI data on SVD markers, while another qualified neurologist (Y-KC) conducted the measurement on infarcts. The numbers and volumes of acute and old infarcts were recorded. The details of the scoring method on 'SVD burden' were described earlier (Liang et al., 2017). Briefly, based on a validated scale (Staals et al., 2014), the presence of either lacune, cerebral microbleeds, extensive WMH, or moderate to severe enlarged perivascular spaces was awarded one point. Summing the above four subscores produced a total SVD score ranging from 0 to 4. The inter-rater reliability of the MRI characteristics was determined by comparing 30 randomly selected MRI scans rated separately by YL and Y-KC. The inter-rater kappa coefficient values were 0.76 for lacune; 0.63 for periventricular WMH, 0.62 for deep WMH; 0.78 for cerebral microbleeds, 0.80 for enlarged perivascular spaces; and 0.66 for both acute and old infarcts.

Statistical Analysis

SPSS Statistics, Version 20.0 software (IBM) was used for the statistical analysis. Data are presented as mean \pm standard deviation (SD), median (interquartile range, IQR), or proportions, as appropriate. The normality of the data was tested with the Shapiro-Wilk test. To ascertain the associations between SVD burden and PDS during the 15-month follow-up, generalized estimating equation (GEE) models were constructed by including patients attending both the first (T1) and the last (T3) follow-ups ($n = 563$). The summed and individual SVD scores were entered the multivariate GEE models, adjusting for demographic, clinical and imaging characteristics. The level of significance was set at 0.05 (two-tailed).

TABLE 1 | Comparison of demographic, clinical and MRI characteristics between the study sample and the patients withdrawn from the study.

	Included ($n = 563$)	Withdrawn ($n = 146$)
Demographics		
Age, years, mean \pm SD	67.0 \pm 10.2	67.1 \pm 11.4
Female sex, n (%)	230 (40.9)	52 (35.6)
Education, years, mean \pm SD	6.7 \pm 4.2	6.7 \pm 5.1
Vascular risk factors, n (%)		
Current or previous smoker	214 (38.0)	63 (42.9)
Hypertension	364 (64.7)	98 (67.6)
Hyperlipidemia	240 (42.6)	52 (35.9)
Diabetes mellitus	159 (28.2)	44 (30.3)
Previous stroke	56 (9.9)	19 (13.1)
Ischemic heart disease	26 (4.6)	8 (5.5)
Atrial fibrillation	24 (4.3)	16 (11.0)
Stroke characteristics		
NIHSS on admission, median (IQR)	3 (1–5)	3 (1–5)
Volume of acute infarcts, ml, mean \pm SD	3.2 \pm 9.6	3.5 \pm 9.1
Number of acute infarcts, mean \pm SD	1.3 \pm 2.2	1.1 \pm 1.1
Old infarcts, n (%)	114 (20.2)	58 (39.7)
Location of acute infarcts, n (%)		
Cortical	96 (17.1)	31 (21.2)
Subcortical white matter	173 (30.7)	36 (24.7)
Deep	127 (22.6)	31 (21.2)
Infratentorial	80 (14.2)	30 (20.5)
SVD individual markers		
Lacune, n (%)	113 (20.1)	28 (19.2)
Cerebral microbleeds, n (%)	76 (13.5)	29 (19.9)
Perivascular WMH, median (IQR)	1 (1–2)	1 (1–2)
Deep WMH, median (IQR)	1 (1–2)	1 (1–2)
Enlarged perivascular spaces, median (IQR)	1 (1–2)	1 (0–2)
SVD burden		
SVD score, median (IQR)	1 (0–2)	1 (0–2)
SVD score = 0, n (%)	248 (44.0)	57 (39.0)
SVD score = 1, n (%)	134 (23.8)	30 (20.5)
SVD score = 2, n (%)	103 (18.3)	36 (24.7)
SVD score = 3, n (%)	54 (9.6)	15 (10.3)
SVD score = 4, n (%)	24 (4.3)	8 (5.5)
Clinical assessments		
Social support: LSNS, mean \pm SD	28.3 \pm 8.4	28.5 \pm 8.7
PDS, n (%)		
T1	103 (18.3)	20 (13.7)
T2	53 (11.6)	–
T3	69 (12.3)	–
Severity of PDS: GDS, median (IQR)		
T1	2 (1–5)	2 (1–4)
T2	2 (1–4)	–
T3	2 (1–4)	–
Functional independence: BI, mean \pm SD		
T1	19.1 \pm 2.1	19.1 \pm 1.9
T2	19.0 \pm 2.3	–
T3	19.1 \pm 2.1	–
Cognitive function: MMSE, mean \pm SD		
T1	26.9 \pm 3.1	26.5 \pm 3.4

(Continued)

TABLE 1 | Continued

	Included (n = 563)	Withdrawn (n = 146)
T2	27.1 ± 3.2	–
T3	26.7 ± 3.7	–

BI, Barthel Index; GDS, Geriatric Depression Scale; IQR, interquartile range; LSNS, Lubben Social Network Scale; MMSE, Mini-Mental State Examination; MRI, Magnetic Resonance Imaging; NIHSS, National Institutes of Health Stroke Scale; PDS, poststroke depressive symptoms; SD, standard deviation; SVD, small vessel disease; WMH, white matter hyperintensities.

RESULTS

Characteristics of the Study Sample

A total of 563 patients were enrolled at both T1 and T3; 146 (20.5%) patients defaulted at follow-up. **Table 1** lists the characteristics of the study sample and the patients who defaulted. The stroke severity of the study sample was lower than that of patients excluded from the study [NIHSS scores of 3 (1–5) vs. 4 (1–11); $p < 0.001$; data not shown].

The demographic, clinical and imaging characteristics are shown in **Table 1**. The study sample had a mean age of 67.0 ± 10.2 years; 40.9% were female. The stroke severity was mainly mild to moderate with a median NIHSS score of 3 (IQR, 1–5); 44.0% of the sample had no marked SVD (SVD score = 0) while the median SVD score was 1 (IQR, 0–2). Levels of social support (LSNS: 28.3 ± 8.4), functional independence (BI: 19.1 ± 2.1) and cognitive functions (MMSE: 26.9 ± 3.1) were relatively high. The level of PDS was low with a median GDS score of 2 across the three assessments. The frequency of clinically meaningful PDS at T1, T2, and T3 was 18.3% (103), 11.6% (53), and 12.4% (69), respectively.

Association Between SVD and PDS

The results of the multivariate GEE model are shown in **Table 2**. The combined SVD score (OR: 1.30, 95% CI: 1.07–1.58; $p = 0.010$) was a significant predictor of PDS over the 15-month follow-up. None of the four individual SVD markers (lacune: $p = 0.720$; cerebral microbleeds: $p = 0.066$; WMH: $p = 0.430$; enlarged perivascular spaces: $p = 0.369$) was significantly associated with PDS in the multivariate GEE models (data not shown).

Other Predictors of PDS

Other significant predictors of PDS were female sex (OR: 1.80, 95% CI: 1.02–3.17, $p = 0.042$); smoking (OR: 1.75, 95% CI: 1.06–2.90, $p = 0.029$); number of acute infarcts (OR: 1.11, 95% CI: 1.02–1.20, $p = 0.014$); time of assessment at T2 (OR: 0.60, 95% CI: 0.42–0.85, $p = 0.004$) and T3 (OR: 0.61, 95% CI: 0.45–0.82, $p = 0.001$) compared with T1; higher level of social support (OR: 0.96, 95% CI: 0.93–0.98, $p < 0.001$); and functional independence (OR: 0.87, 95% CI: 0.81–0.92, $p < 0.001$) (**Table 2**).

TABLE 2 | Multivariate GEE model on the association between PDS and baseline characteristics.

Significant predictors ^a	OR (95%CI)	p
Time points of assessments		
T1	Reference	
T2	0.60 (0.42–0.85)	0.004
T3	0.61 (0.45–0.82)	0.001
Sex		
Male	Reference	
Female	1.80 (1.02–3.17)	0.042
Smoking		
Non-smoker	Reference	
Current or previous smoker	1.75 (1.06–2.90)	0.029
Social support (LSNS)	0.96 (0.93–0.98)	<0.001
SVD burden (SVD score)		
	1.30 (1.07–1.58)	0.010
Number of acute infarcts	1.11 (1.02–1.20)	0.014
Functional independence (BI)	0.87 (0.81–0.92)	<0.001

BI, Barthel Index; CI, confidence interval; GEE, generalized estimating equation; LSNS, Lubben Social Network Scale; OR, odds ratio; PDS, poststroke depressive symptoms; SVD, small vessel disease.

^aAll the predictors listed here were adjusted for assessment time points, age, sex, social support, vascular risk factors, stroke severity, location, number and volume of acute infarcts, old infarcts, SVD burden, cognitive function, and functional independence.

DISCUSSION

The main finding of this study is that SVD burden is associated with clinically meaningful PDS over a 15-month follow-up period in patients with mild to moderate acute ischemic stroke. To the best of our knowledge, this was the first longitudinal study that examined the relationship between SVD burden and PDS after stroke.

It is of note that the SVD burden rather than any single SVD marker was associated with PDS. Higher SVD burden is associated with risk of depression 3 months after lacunar stroke (Zhang X. et al., 2017). The current study extended this finding by conducting a 15-month follow-up to ascertain the long-term impact of the baseline SVD burden on PDS. All subtypes of ischemic stroke were included because lacunar strokes only account for 25% of ischemic strokes (Kolominsky-Rabas et al., 2001). Total SVD burden has been suggested to be a better predictor of stroke outcomes, including mortality (Song et al., 2017), disability (Arba et al., 2017a), stroke recurrence (Lau et al., 2017), and poor quality of life (Liang et al., 2017), than a single SVD marker. The findings of this study lend further support to the hypothesis that chronic vascular burden in the brain is a critical neuroanatomical factor in depression in older adults (Santos et al., 2009b).

The underlying mechanisms linking SVD burden to poststroke PDS are poorly understood. Functional disability and cognitive impairment may be mediating factors as they confer the highest risk for PSD (Towfighi et al., 2017). SVD lesions reduce the brain's 'reserve' against vascular insults and its plasticity (Mok et al., 2017). Stroke patients with more severe SVD are prone to have larger infarct volume (Helenius et al., 2016), impaired reperfusion (Arba et al., 2017b), and

recurrent stroke (Lau et al., 2017). Consequently, disability and cognitive decline were more likely to occur among stroke survivors with severe SVD burden (Huijts et al., 2013; Arba et al., 2017a). However, after adjustment for acute infarcts, stroke severity, functional independence, and cognition in this study, SVD burden remained a significant predictor of PDS. This suggests that SVD burden might contribute to the development of PDS via other paths. From a network viewpoint, the integrity of both the structural and functional global network is the key prerequisite for recovery from stroke (Lim and Kang, 2015) and emotional modulation (Gong and He, 2015). SVD leads to the disruption of white matter integrity of frontal-subcortical circuit which is thought to be critical in the pathogenesis of depression (van Uden et al., 2015). Reduced global structural network efficiency arising from SVD has recently been shown to be associated with depression (Xie et al., 2017).

The pathomechanism of PSD is not yet clarified. In addition to the 'vascular depression hypothesis' as mentioned above, evidence is emerging to support the notion that inflammation could also play a role in the development of PSD (Popa-Wagner et al., 2014; Becker, 2016). Higher level of inflammatory markers was recently found to be predictor of PSD (Cheng et al., 2017) and anti-inflammatory treatment has shown promising results in reducing risk of PSD (Wium-Andersen et al., 2017). Furthermore, SVD burden was found to be associated with systemic lupus erythematosus, a systemic inflammatory disease (Wiseman et al., 2016). Enlarged perivascular spaces, one of the SVD markers, was proposed as an MRI marker of neuroinflammatory activity and was associated with lesions in multiple sclerosis (Wuerfel et al., 2008). The above findings suggest that SVD burden could be an epiphenomenon of the accumulating inflammatory processes.

Medical comorbidities are the 'silent contributors' to stroke outcomes (Sandu et al., 2015). Intracranial atherosclerosis and diabetes were recently reported to be risk factors of early-onset and late-onset PSD, respectively (Chen et al., 2016; Zhang Y. et al., 2017); smoking, as also demonstrated in this study, was associated with PDS. It is suggested that these comorbidities join SVD to promote a chronic proinflammatory state and fuel acute neuroinflammation in the event of acute stroke (Di Napoli et al., 2012; Sandu et al., 2015). The aggravated neuroinflammation, in turn, causes the downstream neurodegeneration, neuronal apoptosis, and impaired neuroplasticity (Leonard and Maes, 2012), disrupt the depression-related neurocircuits (Loubinoux et al., 2012), and eventually result in depressive symptoms or other unfavorable outcomes (Di Napoli et al., 2012). Taken together, PSD or PDS could be viewed as a consequence of a complex interplay between stroke, pre-stroke comorbidities, and neuroinflammation (Sandu et al., 2015).

Other predictors of PDS in the present study were female sex, lower levels of functional independence and social support, smoking, the time of the assessment, and the number of acute infarcts. Female sex, lower level of functional independence and social support have been consistently reported as risk factors of PSD (Towfighi et al., 2017). Smoking also increases the risk of PSD in minor stroke (Shi et al., 2015).

In this study, the risk of PDS was decreasing during the 15-month follow-up. A one-year follow-up study reported that the prevalence of depression decreased from 18.8% at 1 month to 14.6% (Bour et al., 2010). A similar decline in PSD from 6.7% at 3 months to 5.1% at 1 year has also been reported (Kulkantrakorn and Jirapramukpitak, 2007). The number of acute infarcts predicted PDS in the present study, which is consistent with the finding that more severe lacunar infarcts are associated with PSD (Santos et al., 2009a; Wu et al., 2014).

The major strengths of this study are its prospective design, large sample size and its relatively long 15-month follow-up after the index stroke. The first major limitation is that this study mainly included mild to moderate stroke patients. This selection bias reduces the generalizability of the results, including the relationship of SVD burden and PDS in acute ischemic stroke. Second, the lack of comprehensive clinical assessment of depression limits the clinical implications of the study, although PDS also have detrimental effects on stroke outcomes (Towfighi et al., 2017). Nevertheless, the GDS is a reliable tool to detect PDS in this stroke population and has been shown to be useful for screening PDS in Hong Kong elderly stroke patients with good sensitivity and specificity (Tang et al., 2004a,b). A systemic review has also confirmed that the GDS could detect milder PDS in the elderly (Burton and Tyson, 2015). The third limitation is the lack of etiological differentiation of ischemic stroke due to the limited clinical information. Hence, the study could not clarify the differential impact of SVD burden on PDS with respect to the subtypes of acute ischemic stroke. Fourth, about a quarter of eligible patients were lost to follow-up. To offset this limitation, GEE analysis was used to handle missing data. Fifth, this was an observational study without comprehensive investigation of the inflammatory markers or comorbidities. Moreover, we are aware that ideally the assessment of imaging data should be conducted by two raters despite of moderate to good interrater agreements, however, this study failed to provide confirming assessments by a second rater due to the limited manpower. Finally, the 15-month follow-up could not draw conclusions about the association between SVD burden and longer-term clinical course of PDS.

CONCLUSION

SVD burden was associated with clinically significant PDS over a 15-month follow-up in patients with mild to moderate acute ischemic stroke. The findings lend further support to the hypothesis of vascular depression suggesting that accumulating microvascular lesions are critical for the development of depressive symptoms. Measurement of SVD burden might facilitate the early identification of patients at risk of PDS after mild to moderate acute ischemic stroke. Multimodal imaging studies from a network perspective would be useful to clarify the underlying mechanisms linking SVD to PDS. Investigations of the impact of SVD burden on the effectiveness of antidepressant treatment in PDS are also warranted.

AUTHORS CONTRIBUTIONS

W-KT and YL made substantial contributions to the conception and design of the study. YL, Y-KC, VC-TM, D-FW, and WC-WC made substantial contributions to the acquisition of data. YL and Y-KC made substantial contributions to the analysis of data. VM, D-FW, WC-WC, GSU, H-JK, and W-KT contributed to the interpretations of data. YL drafted the first version of the manuscript. All the authors revised the draft for intellectual content, gave their final approval of the final version for publication, and agreed to be accountable for all aspects of the work in ensuring

that questions related to the accuracy or integrity of any part of the study are appropriately investigated and resolved.

FUNDING

This work was supported by a grant from the National Natural Science Foundation of China (Grant Reference Number: 81371460) and a General Research Fund Grant from the University Grants Committee of Hong Kong (Grant Reference Number: 473712).

REFERENCES

- Arba, F., Inzitari, D., Ali, M., Warach, S., Luby, M., and Lees, K. (2017a). Small vessel disease and clinical outcomes after IV rt-PA treatment. *Acta Neurol. Scand.* 136, 72–77. doi: 10.1111/ane.12745
- Arba, F., Mair, G., Carpenter, T., Sakka, E., Sandercock, P., Lindley, R. I., et al. (2017b). Cerebral white matter hypoperfusion increases with small-vessel disease burden. Data from the third international stroke trial. *J. Stroke Cerebrovasc. Dis.* 26, 1506–1513. doi: 10.1016/j.jstrokecerebrovasdis.2017.03.002
- Barthel, D. (1965). Functional evaluation: the Barthel Index. *Md. State Med. J.* 14, 16–65.
- Becker, K. J. (2016). Inflammation and the silent sequelae of stroke. *Neurotherapeutics* 13, 801–810. doi: 10.1007/s13311-016-0451-5
- Bour, A., Rasquin, S., Aben, I., Boreas, A., Limburg, M., and Verhey, F. (2010). A one-year follow-up study into the course of depression after stroke. *J. Nutr. Health Aging* 14, 488–493. doi: 10.1007/s12603-010-0033-x
- Brott, T., Adams, H. P. Jr., Olinger, C. P., Marler, J. R., Barsan, W. G., Biller, J., et al. (1989). Measurements of acute cerebral infarction: a clinical examination scale. *Stroke* 20, 864–870. doi: 10.1161/01.STR.20.7.864
- Burton, L. J., and Tyson, S. (2015). Screening for mood disorders after stroke: a systematic review of psychometric properties and clinical utility. *Psychol. Med.* 45, 29–49. doi: 10.1017/s0033291714000336
- Carson, A. J., MacHale, S., Allen, K., Lawrie, S. M., Dennis, M., House, A., et al. (2000). Depression after stroke and lesion location: a systematic review. *Lancet* 356, 122–126. doi: 10.1016/s0140-6736(00)02448-x
- Chen, Y. K., Qu, J. F., Xiao, W. M., Li, W. Y., Li, W., Fang, X. W., et al. (2016). Intracranial atherosclerosis and poststroke depression in Chinese patients with ischemic stroke. *J. Stroke Cerebrovasc. Dis.* 25, 998–1004. doi: 10.1016/j.jstrokecerebrovasdis.2015.12.038
- Cheng, L. S., Tu, W. J., Shen, Y., Zhang, L. J., and Ji, K. (2017). Combination of high-sensitivity C-reactive protein and homocysteine predicts the post-stroke depression in patients with ischemic stroke. *Mol. Neurobiol.* doi: 10.1007/s12035-017-0549-8 [Epub ahead of print].
- Chiu, H. F., Lee, H., Chung, W., and Kwong, P. (1994). Reliability and validity of the Cantonese version of mini-mental state examination—a preliminary study. *J. Hong Kong Coll. Psychiatry* 4(Suppl. 2), 25–28.
- Di Napoli, M., Godoy, D. A., Campi, V., Masotti, L., Smith, C. J., Parry Jones, A. R., et al. (2012). C-reactive protein in intracerebral hemorrhage: time course, tissue localization, and prognosis. *Neurology* 79, 690–699. doi: 10.1212/WNL.0b013e318264e3be
- Gong, Q., and He, Y. (2015). Depression, neuroimaging and connectomics: a selective overview. *Biol. Psychiatry* 77, 223–235. doi: 10.1016/j.biopsych.2014.08.009
- Helenius, J., Mayasi, Y., and Henninger, N. (2016). White matter hyperintensity lesion burden is associated with the infarct volume and 90-day outcome in small subcortical infarcts. *Acta Neurol. Scand.* 135, 585–592. doi: 10.1111/ane.12670
- Huijts, M., Duits, A., Van Oostenbrugge, R. J., Kroon, A. A., De Leeuw, P. W., and Staals, J. (2013). Accumulation of MRI markers of cerebral small vessel disease is associated with decreased cognitive function: a study in first-ever lacunar stroke and hypertensive patients. *Front. Aging Neurosci.* 5:72. doi: 10.3389/fnagi.2013.00072
- Kolominsky-Rabas, P. L., Weber, M., Gefeller, O., Neundoerfer, B., and Heuschmann, P. U. (2001). Epidemiology of ischemic stroke subtypes according to TOAST criteria. *Stroke* 32, 2735–2740. doi: 10.1161/hs1201.100209
- Kulkantrakorn, K., and Jirapramukpitak, T. (2007). A prospective study in one year cumulative incidence of depression after ischemic stroke and Parkinson's disease: a preliminary study. *J. Neurol. Sci.* 263, 165–168. doi: 10.1016/j.jns.2007.07.014
- Lau, K. K., Li, L., Schulz, U., Simoni, M., Chan, K. H., Ho, S. L., et al. (2017). Total small vessel disease score and risk of recurrent stroke validation in 2 large cohorts. *Neurology* 88, 2260–2267. doi: 10.1212/WNL.0000000000004042
- Leonard, B., and Maes, M. (2012). Mechanistic explanations how cell-mediated immune activation, inflammation and oxidative and nitrosative stress pathways and their sequels and concomitants play a role in the pathophysiology of unipolar depression. *Neurosci. Biobehav. Rev.* 36, 764–785. doi: 10.1016/j.neubiorev.2011.12.005
- Liang, Y., Chen, Y.-K., Deng, M., Mok, V. C. T., Wang, D.-F., Ungvari, G. S., et al. (2017). Association of cerebral small vessel disease burden and health-related quality of life after acute ischemic stroke. *Front. Aging Neurosci.* 9:372. doi: 10.3389/fnagi.2017.00372
- Lim, J. S., and Kang, D. W. (2015). Stroke connectome and its implications for cognitive and behavioral sequela of stroke. *J. Stroke* 17, 256–267. doi: 10.5853/jos.2015.17.3.256
- Loubinoux, I., Kronenberg, G., Endres, M., Schumann-Bard, P., Freret, T., Filipkowski, R. K., et al. (2012). Post-stroke depression: mechanisms, translation and therapy. *J. Cell. Mol. Med.* 16, 1961–1969. doi: 10.1111/j.1582-4934.2012.01555.x
- Lubben, J. E. (1988). Assessing social networks among elderly populations. *Fam. Commun. Health* 11, 42–52. doi: 10.1097/00003727-198811000-00008
- Mok, V. C., Lam, B. Y., Wong, A., Ko, H., Markus, H. S., and Wong, L. K. (2017). Early-onset and delayed-onset post-stroke dementia - revisiting the mechanisms. *Nat. Rev. Neurol.* 13, 148–159. doi: 10.1038/nrneuro.2017.16
- Popa-Wagner, A., Buga, A. M., Tica, A. A., and Albu, C. V. (2014). Perfusion deficits, inflammation and aging precipitate depressive behaviour. *Biogerontology* 15, 439–448. doi: 10.1007/s10522-014-9516-1
- Sandu, R. E., Buga, A. M., Uzoni, A., Petcu, E. B., and Popa-Wagner, A. (2015). Neuroinflammation and comorbidities are frequently ignored factors in CNS pathology. *Neural Regen. Res.* 10, 1349–1355. doi: 10.4103/1673-5374.165208
- Santos, M., Gold, G., Kovari, E., Herrmann, F. R., Bozikas, V. P., Bouras, C., et al. (2009a). Differential impact of lacunes and microvascular lesions on poststroke depression. *Stroke* 40, 3557–3562. doi: 10.1161/strokeaha.109.548545
- Santos, M., Kovari, E., Hof, P. R., Gold, G., Bouras, C., and Giannakopoulos, P. (2009b). The impact of vascular burden on late-life depression. *Brain. Res. Rev.* 62, 19–32. doi: 10.1016/j.brainresrev.2009.08.003
- Shi, Y., Xiang, Y., Yang, Y., Zhang, N., Wang, S., Ungvari, G. S., et al. (2015). Depression after minor stroke: prevalence and predictors. *J. Psychosom. Res.* 79, 143–147. doi: 10.1016/j.jpsychores.2015.03.012
- Song, T. J., Kim, J., Song, D., Yoo, J., Lee, H. S., Kim, Y. J., et al. (2017). Total cerebral small-vessel disease score is associated with mortality during follow-up

- after acute ischemic stroke. *J. Clin. Neurol.* 13, 187–195. doi: 10.3988/jcn.2017.13.2.187
- Staals, J., Makin, S. D., Doubal, F. N., Dennis, M. S., and Wardlaw, J. M. (2014). Stroke subtype, vascular risk factors, and total MRI brain small-vessel disease burden. *Neurology* 83, 1228–1234. doi: 10.1212/WNL.0000000000000837
- Stern, R. A., and Bachman, D. L. (1991). Depressive symptoms following stroke. *Am. J. Psychiatry* 148, 351–356. doi: 10.1176/ajp.148.3.351
- Tang, W. K., Chan, S. S., Chiu, H. F., Wong, K. S., Kwok, T. C., Mok, V., et al. (2004a). Can the geriatric depression scale detect post-stroke depression in Chinese elderly? *J. Affect. Disord.* 81, 153–156. doi: 10.1016/S0165-0327(03)00163-0
- Tang, W. K., Chen, Y., Liang, H., Chu, W. C., Mok, V. C., Ungvari, G. S., et al. (2014). Cerebral microbleeds as a predictor of 1-year outcome of poststroke depression. *Stroke* 45, 77–81. doi: 10.1161/STROKEAHA.113.002686
- Tang, W. K., Ungvari, G. S., Chiu, H. F., and Sze, K. H. (2004b). Detecting depression in Chinese stroke patients: a pilot study comparing four screening instruments. *Int. J. Psychiatry. Med.* 34, 155–163. doi: 10.2190/9YJ9-NNXA-RJCY-2DYP
- Towfighi, A., Ovbiagele, B., El Husseini, N., Hackett, M. L., Jorge, R. E., Kissela, B. M., et al. (2017). Post-stroke depression: a scientific statement for healthcare professionals from the American heart association/American stroke association. *Stroke* 48, e30–e43. doi: 10.1161/STR.0000000000000113
- Uiterwijk, R., van Oostenbrugge, R. J., Huijts, M., De Leeuw, P. W., Kroon, A. A., and Staals, J. (2016). Total cerebral small vessel disease MRI score is associated with cognitive decline in executive function in patients with hypertension. *Front. Aging Neurosci.* 8:301. doi: 10.3389/fnagi.2016.00301
- van Uden, I. W., Tuladhar, A. M., de Laat, K. F., van Norden, A. G., Norris, D. G., van Dijk, E. J., et al. (2015). White matter integrity and depressive symptoms in cerebral small vessel disease: the RUN DMC study. *Am. J. Geriatr. Psychiatry* 23, 525–535. doi: 10.1016/j.jagp.2014.07.002
- Wardlaw, J. M., Smith, E. E., Biessels, G. J., Cordonnier, C., Fazekas, F., Frayne, R., et al. (2013). Neuroimaging standards for research into small vessel disease and its contribution to ageing and neurodegeneration. *Lancet Neurol.* 12, 822–838. doi: 10.1016/S1474-4422(13)70124-8
- Wei, N., Yong, W., Li, X., Zhou, Y., Deng, M., Zhu, H., et al. (2015). Post-stroke depression and lesion location: a systematic review. *J. Neurol.* 262, 81–90. doi: 10.1007/s00415-014-7534-1
- Wiseman, S. J., Bastin, M. E., Jardine, C. L., Barclay, G., Hamilton, I. F., Sandeman, E., et al. (2016). Cerebral small vessel disease burden is increased in systemic lupus erythematosus. *Stroke* 47, 2722–2728. doi: 10.1161/STROKEAHA.116.014330
- Wium-Andersen, I. K., Wium-Andersen, M. K., Jorgensen, M. B., and Osler, M. (2017). Anti-inflammatory treatment and risk for depression after first-time stroke in a cohort of 147 487 Danish patients. *J. Psychiatry Neurosci.* 42, 320–330. doi: 10.1503/jpn.160244
- Wu, R. H., Feng, C., Xu, Y., Hua, T., Liu, X. Y., and Fang, M. (2014). Late-onset depression in the absence of stroke: associated with silent brain infarctions, microbleeds and lesion locations. *Int. J. Med. Sci.* 11, 587–592. doi: 10.7150/ijms.8025
- Wuerfel, J., Haertle, M., Waiczies, H., Tysiak, E., Bechmann, I., Wernecke, K. D., et al. (2008). Perivascular spaces—MRI marker of inflammatory activity in the brain? *Brain* 131, 2332–2340. doi: 10.1093/brain/awn171
- Xie, X., Shi, Y., and Zhang, J. (2017). Structural network connectivity impairment and depressive symptoms in cerebral small vessel disease. *J. Affect. Disord.* 220, 8–14. doi: 10.1016/j.jad.2017.05.039
- Zhang, X., Tang, Y., Xie, Y., Ding, C., Xiao, J., Jiang, X., et al. (2017). Total magnetic resonance imaging burden of cerebral small-vessel disease is associated with post-stroke depression in patients with acute lacunar stroke. *Eur. J. Neurol.* 24, 374–380. doi: 10.1111/ene.13213
- Zhang, Y., He, J. R., Liang, H. B., Lu, W. J., Yang, G. Y., Liu, J. R., et al. (2017). Diabetes mellitus is associated with late-onset post-stroke depression. *J. Affect. Disord.* 221, 222–226. doi: 10.1016/j.jad.2017.06.045

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Liang, Chen, Mok, Wang, Ungvari, Chu, Kang and Tang. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.