
Theses

2010

Utilising a Combined Exercise and Counselling Program to Examine the
Relationship Between Emotional Self-Efficacy and Physiological Improvements in
Breast Cancer Survivors

Jena K. Buchan
University of Notre Dame Australia

Follow this and additional works at: <http://researchonline.nd.edu.au/theses>



COMMONWEALTH OF AUSTRALIA
Copyright Regulations 1969

WARNING

The material in this communication may be subject to copyright under the Act. Any further copying or communication of this material
by you may be the subject of copyright protection under the Act.

Do not remove this notice.

Publication Details

Buchan, J. K. (2010). Utilising a Combined Exercise and Counselling Program to Examine the Relationship Between Emotional Self-Efficacy and Physiological Improvements in Breast Cancer Survivors (Master of Health Sciences). University of Notre Dame Australia.
<http://researchonline.nd.edu.au/theses/45>

This dissertation/thesis is brought to you by ResearchOnline@ND. It has been accepted for inclusion in Theses by an authorized administrator of ResearchOnline@ND. For more information, please contact researchonline@nd.edu.au.



References

- Agnes, C. (2003). Public perceptions of counsellors: a survey of knowledge and attitude to counsellors on the Gold Coast (Doctoral dissertation, Bond University, 2003).. Retrieved from <http://epublications.bond.edu.au/theses/agnew>
- Alder, J. & Bitzer, J. (2003). Retrospective evaluation of the treatment for breast cancer: How does the patient's personal experience of the treatment affect later adjustment to the illness? *Archives of Women's Mental Health*, 6, 91-97.
- American Cancer Society [ACS] (2008). *Cancer Facts and Figures 2008*, Atlanta: American Cancer Society.
- American College of Sports Medicine [ACSM] (2006). *ACSM's guidelines for exercise testing and prescription (7th ed.)*. Philadelphia: Lippincott, Williams & Wilkins.

- Australian Institute of Health and Welfare [AIHW] & Australasian Association of Cancer Registries [AACR] (2008). *Cancer in Australia: an overview, 2008*, Canberra: AIHW.
- Australian Institute of Health and Welfare [AIHW] & National Breast Cancer Centre [NBCC] (2006). *Breast cancer in Australia: an overview, 2006*, Canberra: AIHW.
- Avis, N. E., Crawford, S., & Manuel, J. (2004). Psychosocial problems among younger women with breast cancer. *Psycho-Oncology*, *13*(5), 295-308.
- Bandura, A. (1982). Self-efficacy mechanism in human agency. *American Psychology*, *37*, 112-147.
- Battaglini, C., Bottaro, M., Dennehy, C., Rae, L., Shields, E., Kirk, D., et al. (2007). The effects of an individualized exercise intervention on body composition in breast cancer patients undergoing treatment. *Sao Paulo Medical Journal*, *125*(1), 22-28
- Battaglini, C. L., Dennehy, C. A., Groff, D., Kirk, D., & Anton, P. M. (2006). Complementary therapies in the management of cancer treatment-related symptoms: the individualized prescriptive exercise intervention approach. *Medicina Sportiva*, *10*(2), 49-57.
- Beck, A.T., Ward, C.H., Mendelson, M., Mock, J., & Erbaugh, J. (1961). An inventory for measuring depression. *Archives of General Psychiatry*, *4*, 561-571.
- Berger, A.M. & Farr, L. (1999). The influence of daytime inactivity and nighttime restlessness on cancer-related fatigue. *Oncology Nursing Forum*, *26*, 1663-1671.

- Bezwada, H.P., Granick, M.S., Long, C.D., Moore, J.H., Jr., Lackman, R.L., & Weiss, A.J. (1998). Soft-tissue complications of intra-arterial chemotherapy for extremity sarcomas. *Annals of Plastic Surgery*, 40(4), 382-387.
- Bicego, D., Brown, K., Ruddick, M., Storey, M., Wong, C., & Harris, S.R. (2009). Effects of exercise on quality of life in women living with breast cancer: a systematic review. *The Breast Journal*, 15(1), 45-51.
- Bozoian, S., Rejeski, W.J., & McAuley, E. (1994). Self-efficacy influences feeling states associated with acute exercise. *Journal of Sport and Exercise Psychology*, 16, 326-333.
- Brady, M.J., Cella, D.F., Mo, F., Bonomi, A.E., Tulsy, D.S., Lloyd, S.R., et al. (1997). Reliability and validity of the Functional Assessment of Cancer Therapy-Breast quality-of-life instrument. *Journal of Clinical Oncology*, 15, 974-986.
- Brockstein, B.E., Smiley, C., Al-Sadir, J., & Williams, S.F. (2000). Cardiac and pulmonary toxicity in patients undergoing high-dose chemotherapy for lymphoma and breast cancer: prognostic factors. *Bone Marrow Transplant*, 25, 885-894.
- Bulsara, C., Chan, A., & Styles, I. (2008). *The healthcare team as a resource for empowering breast cancer patients*. Poster presented at GP and PHC Research Conference. Retrieved October 27, 2009, from <http://www.phcris.org.au/roar/profiles.php?elibid=836>
- Byar, K. L., Berger, A. M., Bakken, S. L., & Cetak, M. A. (2006). Impact of adjuvant breast cancer chemotherapy on fatigue, other symptoms, and quality of life. *Oncology Nursing Forum*, 33(1), E18-E26., 8, 1327-1334.

- Camoriano, J.K., Loprinzi, C.I., Ingle, J.N., Themeau, T.M., Krook, J.E., & Veeder, M.H. (1990). Weight change in women treated with adjuvant therapy or observed following mastectomy for node-positive breast cancer. *Journal of Clinical Oncology*, 8, 1327-1334.
- Campbell, A., Mutrie, N., White, F., McGuire, F., & Kearney, N. (2005). A pilot study of a supervised group exercise programme as a rehabilitation treatment for women with breast cancer receiving adjuvant treatment. *European Journal of Oncology Nursing*, 9(1), 56-63.
- Canadian Society for Exercise Physiology [CSEP] (2003). *The Canadian physical activity, fitness and lifestyle approach: CSEP-Health and Fitness Program's Health-Related Appraisal and Counselling Strategy (3rd ed.)*. Ottawa, ON: Canadian Society for Exercise Physiology.
- Carruthers, C. & Hood, C.D. (2004). The power of the positive: leisure and well-being. *Therapeutic Recreation Journal*, 38(2), 225-245.
- Cella, D.F., Tulsky, D.S., Gray, G., Sarafian, B., Linn, E., Bonomi, A., et al. (1993). The Functional Assessment of Cancer Therapy scale: development and validation of the general measure. *Journal of Clinical Oncology*, 11, 570-579.
- Cheney, C. L., Mahloch, J., & Freeny, P. (1997). Computerized tomography assessment of women with weight changes associated with adjuvant treatment for breast cancer. *American Journal of Clinical Nutrition*, 66, 141-146.
- Cimprich, B. (1992). Attentional fatigue following breast cancer surgery. *Research in Nursing & Health*, 15, 199-207.
- Classen, C., Butler, L.D., Koopman, C., Miller, E., DiMiceli, S., Giese-Davis, J., et al. (2001). Supportive-expressive group therapy and distress in patients with metastatic breast cancer. *Archives of General Psychiatry*, 58, 494-501.

- Classen, C., Koopman, C., Angell, K., & Spiegel, D. (1996). Coping styles associated with psychological adjustment to advanced breast cancer. *Health Psychology, 15*, 434-437.
- Classen, C.C., Kraemer, H.C., Blasey, C., Giese-Davis, J., Koopman, C., Palesh, O.G., et al. (2007). Supportive-expressive group therapy for primary breast cancer patients: a randomized prospective multicenter trial. *Psycho-Oncology, 17*(5), 438-447.
- Cohen, Jacob (1988). *Statistical Power Analysis for the Behavioural Sciences* (2nd ed). Lawrence Erlbaum Associates.
- Collie, K., Wong, P., Tilston, J., Butler, L.D., Turner-Cobb, J., Kreshka, M.A., et al. (2005). Self-efficacy, coping, and difficulties interacting with health care professionals among women living with breast cancer in rural communities. *Psycho-Oncology, 14*, 901-912.
- Costa, L. J. M., Varella, P. C. S., & Giglio, A. d. (2002). Weight changes during chemotherapy for breast cancer. *Sao Paulo Medical Journal, 120*(4), 113-117.
- Courneya, K.S. (2003). Exercise in cancer survivors an overview of research. *Medicine & Science in Sports & Exercise, 35*(11), 1846-1852.
- Courneya, K.S., Mackey, J.R., Bell, G.J., Jones, L.W., Field, C.J., & Fairey, A.S. (2003). Randomized controlled trial of exercise training in postmenopausal breast cancer survivors: cardiopulmonary and quality of life outcomes. *Journal of Clinical Oncology, 21*(9), 1651-1652.
- Courneya, K. S., Segal, R. J., Gelmon, K., Reid, R. D., Mackey, J. R., Friedenreich, C. M., et al. (2007a). Six-month follow-up of patient-rated outcomes in a randomized controlled trial of exercise training during breast cancer

chemotherapy. *Cancer Epidemiology, Biomarkers, and Prevention*, 16(12), 2572-2578.

Courneya, K. S., Segal, R. J., Mackey, J. R., Gelmon, K., Reid, R. D., Friedenreich, C. M., et al. (2007b). Effects of aerobic and resistance exercise in breast cancer patients receiving adjuvant chemotherapy: a multicenter randomized controlled trial. *Journal of Clinical Oncology*, 25(28), 4396-4404.

Crowley, S.A. (2003). The effect of a structured exercise program on fatigue, strength, endurance, physical self-efficacy, and functional wellness in women with early stage breast cancer (Doctoral dissertation, University of Michigan, 2003). Retrieved October 28, 2009, from CINAHL Plus database.

Cunningham, A.J., Lockwood, G.A., & Cunningham, J.A. (1991). A relationship between perceived self-efficacy and quality of life in cancer patients. *Patient Education and Counselling*, 17, 71-78.

Daley, A. J., Crank, H., Saxton, J. M., Mutrie, N., Coleman, R., & Roalfe, A. (2007). Randomized trial of exercise therapy in women treated for breast cancer. *Journal of Clinical Oncology*, 25(13), 1713-1721.

Demark-Wahnefried, W., Clipp, E.C., Lipkus, I.M., Lobach, D., Synder, D.C., Sloane, R., et al. (2007). Main outcomes of the FRESH START trial: a sequentially tailored, diet and exercise mailed print intervention among breast and prostate cancer survivors. *Journal of Clinical Oncology*, 25(19), 2709-2718.

Demark-Wahnefried, W., Kenyon, A. J., Eberle, P., Skye, A., & Kraus, W. E. (2002). Preventing sarcopenic obesity among breast cancer patients receiving adjuvant chemotherapy: a feasibility study. *Clinical Exercise Physiology*, 4(1), 44-49.

- Demark-Wahnefried, W., Peterson, B. L., Winer, E. P., Marks, L., Aziz, N., Marcom, P. K., et al. (2001). Changes in weight, body composition, and factors influencing energy balance among premenopausal breast cancer patients receiving adjuvant chemotherapy. *Journal of Clinical Oncology*, 19(9), 2381-2389.
- Derogatis, L.R., Morrow, G.R., Fetting, J., Penman, D., Piasetsky, S., Schmale, A.M., et al. (1983). The prevalence of psychiatric disorders among cancer patients. *Journal of the American Medical Association*, 249, 751-757.
- Devine, M.A. & Dattilo, J. (2000). Expressive Arts and Therapeutic Media. In J. Dattilo (Ed.), *Facilitation techniques in therapeutic recreation* (pp. 133-164). State College, PA: Venture Publishing.
- Di Sipio, T., Harrison, S., Newton, M., Guy, L., Spathonis, K., Eakin, E., et al. (2009, May). *A randomised controlled trial of an exercise intervention-quality of life benefits*. Poster session presented at the 15th UICC Reach to Recovery International Breast Cancer Support Conference, Brisbane, QLD.
- Drouin, J. (2002). Aerobic exercise training effects on physical function, fatigue and mood, immune status, and oxidative stress in subjects undergoing radiation treatment for breast cancer (Doctoral dissertation, Wayne State University). Retrieved October 27, 2009, from CINAHL Plus database.
- Dzewaltowski, D.A., Noble, J.M., & Shaw, J.M. (1990). Physical activity participation: social cognitive theory versus the theories of reasoned action and planned behaviour. *Journal of Sport and Exercise Psychology*, 12, 388-405.
- Engel, J., Kerr, J., Schlesinger-Raab, A., Eckler, R., Sauer, H., & Holzel, D. (2003). Predictors of quality of life of breast cancer patients. *Acta Oncologica*, 42, 710-718.

- Eversley, R., Estrin, D., Dibble, S. Wardlaw, L., Pedrosa, M., & Favila-Penney, W. (2005). Post-treatment symptoms among ethnic minority breast cancer survivors. *Oncology Nursing Forum*, 32, 250-256.
- Falagas, M. E., Zarkadoulia, E. A., Ioannidou, E. N., Peppas, G., Christodoulou, C., & Rafailidis, P. I. (2007). The effect of psychosocial factors on breast cancer outcome: a systematic review. *Breast Cancer Research*, 9(4), R44.
- Fleming, M. P., & Kleinbart, E. (2001). Breast cancer and sexuality. *Journal of Sex Education and Therapy*, 26(3), 215-224.
- Freedman, R. J., Aziz, N., Albanes, D., Hartman, T., Danforth, D., Hill, S., et al. (2004). Weight and body composition changes during and after adjuvant chemotherapy in women with breast cancer. *Journal of Clinical Endocrinology and Metabolism*, 89(5), 2248-2253.
- Ganz, P.A., Coscarelli, A., Fred, C., Kahn, B., Polinsky, M.I., & Petersen, L. (1996). Breast cancer survivors: psychosocial concerns and quality of life. *Breast Cancer Research and Treatment*, 38, 183-199.
- Ganz, P.A., Kwan, L., Stanton, A. L., Krupnick, J. L., Rowland, J. H., Meyerowitz, B. E., et al. (2004). Quality of life at the end of primary treatment of breast cancer: first results from the Moving Beyond Cancer randomized trial. *Journal of the National Cancer Institute*, 96(5), 376-387.
- Garreau, J. R., DeLaMelena, T., Walts, D., Karamlou, K., & Johnson, N. (2006). Side effects of aromatase inhibitors versus tamoxifen: the patients' perspective. *The American Journal of Surgery*, 192, 3.
- Giese-Davis, J., Bliss-Isberg, C., Carson, K., Star, P., Donaghy, J., Cordova, M., et al. (2006). The effect of peer counselling on quality of life following diagnosis of breast cancer: an observational study. *Psycho-Oncology*, 15, 1014-1022.

- Giese-Davis, J., Koopman, C., Butler, L.D., Classen, C., Cordova, M., Fobair, et al. (2002). Change in emotion-regulation strategy for women with metastatic breast cancer following supportive-expressive group therapy. *Journal of Consulting and Clinical Psychology, 70*(4), 916-925.
- Giese-Davis, J., Koopman, C., Butler, L. D., Joss, J., Classen, C., Roberts, J., et al. (2004). The Stanford Emotional Self-Efficacy Scale--Cancer: Reliability, validity, and generalisability. In I. Nyklíček, L. Temoshok, & A. Vingerhoets (Eds.). *Emotional Expression and Health: Advances in Theory, Assessment and Clinical Applications* (pp. 204-222). Hove, UK & New York: Brunner-Routledge.
- Giese-Davis, J. & Spiegel, D. (2003). Emotional expression and cancer progression. In Davidson, R.J., Scherer, K.R., & Hill Goldsmith, H. (Eds.), *Handbook of affective sciences* (pp. 1053-1082. Oxford: Oxford University Press.
- Girgis, A., Boyes, A., & Hansen, V. (2008). What are the unmet needs of breast cancer survivors 5-6 years post-diagnosis? *Clinical Updates in Breast Cancer, 3*(2), 15-17.
- Golding, L.A. (1989). *YMCA fitness testing and assessment manual*. Champaign, IL: Human Kinetics.
- Goodwin, P., Esplen, M.J., Butler, K., Winocur, J., Pritchard, K., Brazel, S., et al. (1998). Multidisciplinary weight management in locoregional breast cancer: results of a phase II study. *Breast Cancer Research and Prevention, 48*(1), 53-64.
- Goodwin, P.J., Ennis, M., Pritchard, K.I., Trudeau, M.E., Koo, J., Madarnas, Y., et al. (2002). Fasting insulin and outcome in early stage breast cancer: results of a prospective cohort study. *Journal of Clinical Oncology, 20*(1), 42-51.

- Greenberg, L.S. & Foerster, F.S. (1996). Task analysis exemplified: the process of resolving unfinished business. *Journal of Consulting & Clinical Psychology*, *61*, 78-84.
- Groff, D., Battaglini, C., O'Keefe, C., Edwards, C., & Peppercorn, J. (2007). Lessons from Survivors: The Role of Recreation Therapy in Facilitating Spirituality and Well-Being. Retrieved March 8, 2009, from <http://www.heartmath.org/research/rd-lessons-from-survivors-the-role-of-recreation-therapy-in-facilitating-spirituality.html>
- Groff, D. & Dattilo, J. (2000). Adventure Therapy. In J. Dattilo (Ed.), *Facilitation techniques in therapeutic recreation* (pp. 13-39). State College, PA: Venture Publishing.
- Gross, J.J. (1989). Emotional expression in cancer onset and progression. *Social Science in Medicine*, *28*, 1239-1248.
- Gross, J.J. & Levenson, R.W. (1997). Hiding feelings: the acute effects of inhibiting positive and negative emotions. *Journal of Abnormal Psychology*, *106*, 95-103.
- Han, W. T., Collie, K., Koopman, C., Azarow, J., Classen, C., Morrow, G. P., et al. (2005). Breast cancer and problems with medical interactions: relationships with traumatic stress, emotional self-efficacy, and social support. *Psycho-Oncology*, *14*, 318-330.
- Harrison, S., Eakin, E., Newton, M., Guy, L., Di Sipio, T., & Hayes, S. (2009, May). *A randomised controlled trial of an exercise intervention-physical activity benefits*. Poster session presented at the 15th UICC Reach to Recovery International Breast Cancer Support Conference, Brisbane, QLD.
- Hayes, P.A., Sowood, P.J., Belyavin, A., Cohen, J.B., & Smith, F.W. (1988). Subcutaneous fat thickness measured by magnetic resonance imaging,

ultrasound, and callipers. *Medicine and Science in Sports and Exercise*, 20, 303-309.

Hayes, S., Battistutta, D., & Newman, B. (2005). Objective and subjective upper body function six months following diagnosis of breast cancer. *Breast Cancer Research and Treatment*, 94, 1-10.

Hayes, S., Cornish, B., & Newman, B. (2005). Comparison of methods to diagnose lymphoedema among breast cancer survivors: 6-month follow-up. *Breast Cancer Research and Treatment*, 89, 221-226.

Hayes, S. & Eakin, E. (2006). *Exercise for Health: a breast cancer recovery project*. Instructions manual distributed to participants in Exercise for Health program, at Queensland University of Technology.

Heber, D., Ingles, S., Ashley, J. M., Maxwell, M. H., Lyons, R. F., & Elashoff, R. M. (1996). Clinical detection of sarcopenic obesity by bioelectrical impedance analysis. *American Journal of Clinical Nutrition*, 64, 472S-477S.

Helms, R.L., O'Hea, E.L., & Corso, M. (2008). Body image issues in women with breast cancer. *Psychology, Health & Medicine*, 13(3), 313-325.

Heyward, V. & Wagner, D. (2004). *Applied body composition assessment* (2nd ed.). Champaign, IL: Human Kinetics.

Heyward, V.H. (2006). *Advanced fitness assessment and exercise prescription* (5th ed.). Champaign, IL: Human Kinetics.

Holmes, M. D., Chen, W. Y., Feskanich, D., Kroenke, C. H., & Colditz, G. A. (2005). Physical activity and survival after breast cancer diagnosis. *Journal of the American Medical Association*, 293(20), 2479-2486.

- Hsieh, C.C., Sprod, L.K., Hydock, D.S., Carter, S.D., Hayward, R., & Schneider, C.M. (2008). Effects of a supervised exercise intervention on recovery from treatment regimens in breast cancer survivors. *Oncology Nursing Forum*, 35(6), 909-915.
- Irwin, M.L., Varma, K., Alvarez-Reeves, M., Cadmus, L., Wiley, A., Chung, G.G., et al. (2008). Exercise on insulin and insulin-like growth factors in breast cancer survivors: the Yale Exercise and Survivorship Study. *Cancer Epidemiology, Biomarkers & Prevention*, 18, 306-313.
- Jackson, A.S., Pollock, M.I., & Ward, A. (1980). Generalized equations for predicting body density of women. *Medicine and Science in Sports and Exercise*, 12(3), 175-181.
- Jackson, A.W. & Baker, A.A. (1986). The relationship of the sit and reach test to criterion measures of hamstring and back flexibility in young females. *Research Quarterly for Exercise and Sport*, 57(3), 183-186.
- Jacobsen, P.B., Hann, D.M., Azzarello, L.M., Horton, J., Balducci, L., & Lyman, G.H. (1999). Fatigue in women receiving adjuvant chemotherapy for breast cancer: characteristics, course, and correlates. *Journal of Pain and Symptom Management*, 18, 233-242.
- Jensen, M.R. (1987). Psychobiological factors predicting the course of breast cancer. *Journal of Personality*, 55, 317-342.
- Karvinen, K.H., Courneya, K.S., Campbell, K.L., Pearcey, R.G., Dundas, G., Capstick, V., et al. (2007). Correlates of exercise motivation and behaviour in a population-based sample of endometrial cancer survivors: an application of the Theory of Planned Behaviour. *International Journal of Behavioural Nutrition and Physical Activity*, 4, 21-30.

- Khan, M.A., Sehgal, A., Mitra, B., Agarwal, P.N., Lal, P., & Malik, V.K. (2000). Psychobehavioural impact of mastectomy. *Journal of the Indian Academy of Applied Psychology*, 26(1-2), 65-71.
- Kim, C., Kang, D., & Park, J. (2009). A meta-analysis of aerobic exercise interventions for women with breast cancer. *Western Journal of Nursing Research*, 31(4), 437-461.
- Kirshbaum, M.N. (2006). A review of the benefits of whole body exercise during and after treatment for breast cancer. *Journal of Clinical Nursing*, 16(1), 104-121.
- Knobf, M. T., Insogna, K., DiPietro, L., Fennie, K., & Thompson, A. S. (2008). An aerobic weight-loaded pilot exercise intervention for breast cancer survivors: bone remodeling and body composition outcomes. *Biological Research for Nursing*, 10(1), 34-43.
- Koopman, C., Angell, K., Turner-Cobb, J.M., Kreshka, M.A., Donnelly, P., McCoy, R., et al. (2001). Distress, coping, and social support among rural women recently diagnosed with primary breast cancer. *The Breast Journal*, 7(1), 25-33.
- Kroenke, C. H., Chen, W. Y., Rosner, B., & Holmes, M. D. (2005). Weight, weight gain, and survival after breast cancer diagnosis. *Journal of Clinical Oncology*, 23(7), 1370-1378.
- Kumar, N.B., Allen, K., Cantor, A., Cox, C.E., Greenberg, A., Shah, S., et al. (1997). Weight gain associated with adjuvant tamoxifen therapy in stage I and II breast cancer, fact or artifact? *Breast Cancer Research and Treatment*, 44, 135-143.
- Lermen, J., Bruce, R.A., Sivarajan, E., Pettet, G., & Trimble, S. (1976). Low-level dynamic exercises for earlier cardiac rehabilitation: aerobic and

hemodynamic responses. *Archives of Physical Medicine and Rehabilitation*, 57, 355-360.

Levine, E.G., Raczynski, J.M., & Carpenter, J.T. (1991). Weight gain with breast cancer adjuvant treatment. *Cancer*, 67(7), 1954-1959.

Lieberman, M.A., Golant, M., Giese-Davis, J., Winzlenberg, A., Benjamin, H., Humphreys, K., et al. (2003). Electronic support groups for breast carcinoma: a clinical trial of effectiveness. *Cancer*, 97, 920-925.

Lohman, T.G., Pollock, M.L., Slaughter, M.H., Brandon, L.J., & Boileau, R.A. (1984). Methodological factors and the prediction of body fat in female athletes. *Medicine and Science in Sports and Exercise*, 16, 92-96.

Lönnqvist, J.E., Paunonen, S., Verkasalo, M., Leikas, S., Tuulio-Henriksson, A., & Lönnqvist, J. (2007). Personality characteristics of research volunteers, *European Journal of Personality*, 21(8), 1017-1030.

MacVicar, M., Winningham, M., & Nickel, J. (1989). Effects of aerobic interval training on cancer patients' function capacity. *Nursing Research*, 38, 348-351.

Manuel, J. C., Burwell, S. R., Crawford, S. L., Lawrence, R. H., Farmer, D. F., Hege, A., et al. (2007). Younger women's perceptions of coping with breast cancer. *Cancer Nursing*, 30(2), 10.

Markes, M., Brockow, T., & Resch, K. L. (2006). Exercise for women receiving adjuvant therapy for breast cancer, *Cochrane Database of Systematic Reviews* 2006, 4.

Matthews, C.E., Wilcox, S., Hanby, C.L., Der Ananian, C., Heiney, S.P., Gebretsadik, T., et al. (2007). Evaluation of a 12-week home-based walking intervention for breast cancer survivors. *Supportive Care in Cancer*, 15(2), 203-211.

- McAuley, E., Lox, C., & Duncan, T.E. (1993). Long-term maintenance of exercise, self-efficacy, and physiological change in older adults. *Journal of Gerontology, 48*(4), 218-224.
- McInnes, J. A., & Knobf, M. T. (2001). Weight gain and quality of life in women treated with adjuvant chemotherapy for early-stage breast cancer. *Oncology Nursing Forum, 28*(4), 675-684.
- McNeely, M.L., Campbell, K.L., Rowe, B.H., Klassen, T.P., Mackey, J.R., & Courneya, K.S. (2006). Effects of exercise on breast cancer patients and survivors: a systematic review and meta-analysis. *Canadian Medical Association Journal, 175*(1), 34-41.
- Mills, M., Black, A., Campbell, A., Cardwell, C.R., Galway, K., & Donnelly, M. (2009). Multidimensional rehabilitation programmes for adult cancer survivors (Protocol). *Cochrane Database of Systematic Reviews 2009, 2*.
- Milne, H. M., Gordon, S., Guilfoyle, A., Wallman, K. E., & Courneya, K. S. (2007). Association between physical activity and quality of life among Western Australian breast cancer survivors. *Psycho-Oncology, 16*(12), 10.
- Milne, H. M., Wallman, K. E., Gordon, S., & Courneya, K. S. (2008). Effects of a combined aerobic and resistance exercise program in breast cancer survivors: a randomized controlled trial. *Breast Cancer Research and Treatment, 108*, 279-288.
- Morasso, G., Costantini, M., Viterbori, P., Bonci, F., Del Mastro, L., Musso, M., et al. (2001). Predicting mood disorders in breast cancer patients. *European Journal of Cancer, 37*(2), 216-223.
- Morris, J. & Ingham, R. (1988). Choice of surgery for early breast cancer: Psychosocial considerations. *Social Science and Medicine, 27*, 1257-1262.

- Mosher, C. E., Fuemmeler, B. F., Sloane, R., Kraus, W. E., Lobach, D. F., Snyder, D. C., et al. (2008). Change in self-efficacy partially mediates the effects of the FRESH START intervention on cancer survivors' dietary outcomes. *Psycho-Oncology, 17*, 1014-1023.
- Mutrie, N., Campbell, A. M., Whyte, F., McConnachie, A., Emslie, C., Lee, L., et al. (2007). Benefits of supervised group exercise programme for women being treated for early stage breast cancer: pragmatic randomised controlled trial. *British Medical Journal, 334*, 7.
- Mutrie, N., Lauglan, C., Campbell, M., Marsed, E., & McCarron, T. (1997). The transtheoretical model applied to four Scottish populations. *Journal of Sports Science, 15*, 100.
- Newell, S.A., Sanson-Fisher, R.W., & Savolainen, N.J. (2002). Systematic review of psychological therapies for cancer patients: overview and recommendations for future research. *Journal of the National Cancer Institute, 94*(8), 558-584.
- Nguyen, M., Stewart, R., Banerji, M., Gordon, D., & Kral, J. (2001). Relationships between tamoxifen use, liver fat and body fat distribution in women with breast cancer. *International Journal of Obesity, 25*(2), 3.
- Ohira, T., Schmitz, K.H., Ahmed, R.L., & Yee, D. (2006). Effects of weight training on quality of life in recent breast cancer survivors. *Cancer, 106*(9), 2076-2083.
- Palesh, O.G., Shaffer, T., Larson, J., Edsall, S., Chen, X., Koopman, C., et al. (2006). Emotional self-efficacy, stressful life events, and satisfaction with social support in relation to mood disturbance among women living with breast cancer in rural communities. *The Breast Journal, 12*(2), 123-129.

- Partridge, A. H., Burnstein, H. J., & Winter, E. P. (2001). Side effects of chemotherapy and combined chemohormonal therapy in women with early-stage breast cancer. *Journal of the National Cancer Institute Monographs*, 30, 135-142.
- Pinto, B.M., Clark, M.M., Maruyaman, N.C., & Feder, S.I. (2003). Psychological and fitness changes associated with exercise participation among women with breast cancer. *Psycho-Oncology*, 12, 118-126.
- Pinto, B. M., & Trunzo, J. J. (2004). Body esteem and mood among sedentary and active breast cancer survivors. *Mayo Clinic Proceedings*, 79, 181-186.
- Pinto, B. M., Trunzo, J. J., Reiss, P., & Shiu, S.-Y. (2002). Exercise participation after diagnosis of breast cancer: trends and effects on mood and quality of life. *Psycho-Oncology*, 11, 389-400.
- Piper, B.F., Dibble, S.L., Dodd, M.J., Weiss, M.C., Slaughter, R.E., & Paul, S.M. (1998). The revised Piper Fatigue Scale: psychometric evaluation in women with breast cancer. *Oncology Nursing Forum*, 25, 677-684.
- Rendle, Karen (1997). Survivorship and breast cancer: the psychosocial issues. *Journal of Clinical Nursing*, 6, 403-410.
- Richardson, P. & Handal, P. (1995). The public's perception of psychotherapy and counselling: different views on the effectiveness of psychologists, psychiatrists, and other providers. *Journal of Contemporary Psychotherapy*, 25(4), 367-385.
- Rock, C.L., Flatt, S.W., Newman, V., Caan, B.J., Haan, M.N., Stefanick, M.L., et al. (1999). Factors associated with weight gain in women after diagnosis of breast cancer. *Journal of the American Dietetic Association*, 99(10), 8.

- Rogers, L.Q., Shah, P., Dunnington, G., Greive, A., Shanmugham, A., Dawson, B., et al. (2005). Social cognitive theory and physical activity during breast cancer treatment. *Oncology Nursing Forum*, 32(4), 807-815.
- Rovniak, L.S., Anderson, E.S., Winett, R.A., & Stephens, R.S. (2002). Social cognitive determinants of physical activity in young adults: a prospective structural equation analysis. *Annals of Behavioural Medicine*, 24(2), 149-156.
- Salkind, M.R. (1969). Beck depression inventory in general practice. *Journal of the Royal College of General Practitioners*, 18, 267-271.
- Schain, W. S., d'Angelo, T. M., Dunn, M. E., Lichter, A. S., & Pierce, L. J. (1994). Mastectomy versus conservative surgery and radiation therapy: psychosocial consequences. *Cancer*, 73, 1221-1228.
- Schmid-Büchi, S., Halfens, R.J.G., Dassen, T., & van den Borne, B. (2008). A review of psychosocial needs of breast-cancer patients and their relatives. *Journal of Clinical Nursing*, 17, 2895-2909.
- Schmitz, K. H., Ahmed, R. L., Hannan, P. J., & Yee, D. (2005). Safety and efficacy of weight training in recent breast cancer survivors to alter body composition, insulin, and insulin-like growth factor axis proteins. *Cancer Epidemiology, Biomarkers, and Prevention*, 14(7), 9.
- Schneider, C.M., Dennehy, C.A., & Carter, S.D. (2003). *Exercise and cancer recovery*. Champaign, IL: Human Kinetics.
- Schneider, C.M., Hsieh, C.C., Sprod, L.K., Carter, S.D., & Hayward, R. (2007). Effects of supervised exercise training on cardiopulmonary function and fatigue in breast cancer survivors during and after treatment. *Cancer*, 110(4), 918-925.

- Schwartz, A. L. (2000). Exercise and weight gain in breast cancer patients receiving chemotherapy. *Cancer Practice*, 8(5), 231-237.
- Schwartz, A. L. (2004). Physical activity after a cancer diagnosis: psychosocial outcomes. *Cancer Investigation*, 22(1), 82-92.
- Segal, R., Evans, W., Johnson, D., Smith, J., Colletta, S., Gayton, J., et al. (2001). Structured exercise improves physical functioning in women with stages I and II breast cancer: results of a randomized controlled trial. *Journal of Clinical Oncology*, 19(3), 657-665.
- Sloan, J. A., Cella, D., Frost, M.H., Guyatt, G.H., Sprangers, M.A.G., Symonds, T., et al. (2002). Assessing clinical significance in measuring oncology patient quality of life: introduction to the symposium, content overview, and definition of terms. *Mayo Clinic Proceedings*, 77(4), 367-370.
- Spiegel, D., Butler, L.D., Giese-Davis, J., Koopman, C., Miller, E., DiMiceli, S., et al. (2007). Effects of supportive-expressive group therapy on survival of patients with metastatic breast cancer. *Cancer*, 110(5), 1130-1138.
- Stewart, B. W. & Kleihues P. (Eds) (2003). *World Cancer Report*, Lyon: IARC Press.
- Truong, P.T., Olivotto, I.A., Whelan, T.J., & Levine, M. (2004). Clinical practice guidelines for the care and treatment of breast cancer: 16. Locoregional post-mastectomy radiotherapy. *Canadian Medical Association Journal*, 170(6), 1263-1273.
- Turner, J., Hayes, S., & Reul-Hirche, H. (2004). Improving the physical status and quality of life of women treated for breast cancer: a pilot study of a structured exercise intervention. *Journal of Surgical Oncology*, 86, 141-146.

- Valois, R.F., Umstattd, M.R., Zullig, K.J., & Paxton, R.J. (2008). Physical activity behaviors and emotional self-efficacy: is there a relationship for adolescents? *Journal of School Health, 78*(6), 321-327.
- Visovsky, C. (2006). Muscle strength, body composition, and physical activity in women receiving chemotherapy for breast cancer. *Integrative Cancer Therapies, 5*(3), 183-191.
- Watson, M., Greer, S., Rowden, L., Gorman, C., Roberston, B., Bliss, J.M., et al. (1991). Relationships between emotional control, adjustment to cancer and depression and anxiety in breast cancer patients. *Psychological Medicine, 21*, 51-57.
- Weihl, K.L., Enright, T.M., Simmens, S.J., & Reiss, D. (2000). Negative affectivity, restriction of emotions, and site of metastases predict mortality in recurrent breast cancer. *Journal of Psychosomatic Research, 49*, 59-68.
- Wilmoth, M. C., Coleman, E. A., Smith, S. C., & Davis, C. (2004). Fatigue, weight gain, and altered sexuality in patients with breast cancer: exploration of a symptom cluster. *Oncology Nursing Forum, 31*(6), 1069-1075.
- Winningham, M.L. (1991). Walking program for people with cancer. Getting started. *Cancer Nursing, 14*, 270-276.
- Winningham, M.L., Nail, L.M., Burke, M.B., Brophy, L., Cimprich, B., Jones, L.S., et al. (1994). Fatigue and the cancer experience: the state of the knowledge. *Oncology Nursing Forum, 21*, 23-26.
- Zamboni, M., Armellini, F., Harris, T., Turcato, E., Micciolo, R., Bergamo-Andreis, I.A., et al. (1997). Effects of age on body fat distribution and cardiovascular risk factors in women. *American Journal of Clinical Nutrition, 66*, 111-115.