

---

Theses

---

2018

Assessment of the neuroprotective efficacy of poly-arginine-18 (R18) peptides in a pre-clinical model of perinatal hypoxic-ischaemic encephalopathy (HIE)

Adam Edwards  
*The University of Notre Dame Australia*

Follow this and additional works at: <https://researchonline.nd.edu.au/theses>



COMMONWEALTH OF AUSTRALIA  
Copyright Regulations 1969

WARNING

The material in this communication may be subject to copyright under the Act. Any further copying or communication of this material by you may be the subject of copyright protection under the Act.

Do not remove this notice.

---

Publication Details

Edwards, A. (2018). Assessment of the neuroprotective efficacy of poly-arginine-18 (R18) peptides in a pre-clinical model of perinatal hypoxic-ischaemic encephalopathy (HIE) (Doctor of Philosophy (College of Health Sciences)). University of Notre Dame Australia. <https://researchonline.nd.edu.au/theses/232>

This dissertation/thesis is brought to you by ResearchOnline@ND. It has been accepted for inclusion in Theses by an authorized administrator of ResearchOnline@ND. For more information, please contact [researchonline@nd.edu.au](mailto:researchonline@nd.edu.au).



## 7.5 REFERENCES

- Ashwal, S., Tone, B., Tian, H. R., Chong, S., & Obenaus, A. (2007). Comparison of Two Neonatal Ischemic Injury Models Using Magnetic Resonance Imaging. *Pediatric Research*, *61*(1), 9–14. <https://doi.org/10.1203/01.pdr.0000251612.16069.4b>
- Barks, J. D., Liu, Y., Shangguan, Y., Djuric, Z., Ren, J., & Silverstein, F. S. (2017). Maternal high-fat diet influences outcomes after neonatal hypoxic-ischemic brain injury in rodents. *Journal of Cerebral Blood Flow & Metabolism*, *37*(1), 307–318. <https://doi.org/10.1177/0271678X15624934>
- Committee on Fetus and Newborn, C. O. F. A., Papile, L.-A., Baley, J. E., Benitz, W., Cummings, J., Carlo, W. A., ... Wang, K. S. (2014). Hypothermia and neonatal encephalopathy. *Pediatrics*, *133*(6), 1146–1150. <https://doi.org/10.1542/peds.2014-0899>
- Davidson, J. O., Rout, A. L., Wassink, G., Yuill, C. A., Zhang, F. G., Green, C. R., ... Gunn, A. J. (2015). Non-Additive Effects of Delayed Connexin Hemichannel Blockade and Hypothermia after Cerebral Ischemia in Near-Term Fetal Sheep. *Journal of Cerebral Blood Flow & Metabolism*, *35*(12), 2052–2061. <https://doi.org/10.1038/jcbfm.2015.171>
- Edwards, A. B., Anderton, R. S., Knuckey, N. W., & Meloni, B. P. (2016). Characterisation of neuroprotective efficacy of modified poly-arginine-9 (R9) peptides using a neuronal glutamic acid excitotoxicity model. *Molecular and Cellular Biochemistry*. <https://doi.org/10.1007/s11010-016-2882-z>
- Faulkner, S., Bainbridge, A., Kato, T., Chandrasekaran, M., Kapetanakis, A. B.,

- Hristova, M., ... Robertson, N. J. (2011). Xenon augmented hypothermia reduces early lactate/N-acetylaspartate and cell death in perinatal asphyxia. *Annals of Neurology*, 70(1), 133–150. <https://doi.org/10.1002/ana.22387>
- Hagberg, H., Bona, E., Gilland, E., & Puka-Sundvall, M. (1997). Hypoxia-ischaemia model in the 7-day-old rat: possibilities and shortcomings. *Acta Paediatrica*, 86(S422), 85–88. <https://doi.org/10.1111/j.1651-2227.1997.tb18353.x>
- Hagberg, H., Gilland, E., Diemer, N. H., & Andiné, P. (1994). Hypoxia-ischemia in the neonatal rat brain: histopathology after post-treatment with NMDA and non-NMDA receptor antagonists. *Biology of the Neonate*, 66(4), 205–213. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/7865635>
- Hall, N. J., Eaton, S., Peters, M. J., Hiorns, M. P., Alexander, N., Azzopardi, D. V, & Pierro, A. (2010). Mild controlled hypothermia in preterm neonates with advanced necrotizing enterocolitis. *Pediatrics*, 125(2), e300-8. <https://doi.org/10.1542/peds.2008-3211>
- Jacobs, S. E., Berg, M., Hunt, R., Tarnow-Mordi, W. O., Inder, T. E., & Davis, P. G. (2013). Cooling for newborns with hypoxic ischaemic encephalopathy. *Cochrane Database of Systematic Reviews*. <https://doi.org/10.1002/14651858.CD003311.pub3>
- Liu, X., Dingley, J., Scull-Brown, E., & Thoresen, M. (2015). Adding 5 h delayed xenon to delayed hypothermia treatment improves long-term function in neonatal rats surviving to adulthood. *Pediatric Research*, 77(6), 779–783. <https://doi.org/10.1038/pr.2015.49>
- MacDougall, G., Anderton, R. S., Edwards, A. B., Knuckey, N. W., & Meloni, B. P. (2016). The Neuroprotective Peptide Poly-Arginine-12 (R12) Reduces Cell

Surface Levels of NMDA NR2B Receptor Subunit in Cortical Neurons; Investigation into the Involvement of Endocytic Mechanisms. *Journal of Molecular Neuroscience : MN*. <https://doi.org/10.1007/s12031-016-0861-1>

Meloni, B., Brookes, L., Clark, V., Cross, J., Edwards, A., Anderton, R., ... Knuckey, N. (2015). Poly-arginine and arginine-rich peptides are neuroprotective in stroke models. *J Cereb Blood Flow Metab*. <https://doi.org/10.1038/jcbfm.2015.11>

Meloni, B., Milani, D., Cross, J., Clark, V., Edwards, A., Anderton, R., ... Knuckey, N. (2017). Assessment of the Neuroprotective Effects of Arginine-Rich Protamine Peptides, Poly-Arginine Peptides (R12-Cyclic, R22) and Arginine–Tryptophan-Containing Peptides Following In Vitro Excitotoxicity and/or Permanent Middle Cerebral Artery Occlusion in Rats. *NeuroMolecular Medicine*, 19(2–3), 271–285. <https://doi.org/10.1007/s12017-017-8441-2>

Meloni, B., Milani, D., Edwards, A., Anderton, R., O'Hare Doig, R., Fitzgerald, M., ... Knuckey, N. (2015). Neuroprotective peptides fused to arginine-rich cell penetrating peptides: Neuroprotective mechanism likely mediated by peptide endocytic properties. *Pharmacology & Therapeutics*, 153, 36–54. <https://doi.org/10.1016/j.pharmthera.2015.06.002>

Milani, D., Clark, V., Cross, J., Anderton, R., Knuckey, N., & Meloni, B. (2016). Poly-arginine peptides reduce infarct volume in a permanent middle cerebral artery rat stroke model. *BMC Neuroscience*, 17(1), 19. <https://doi.org/10.1186/s12868-016-0253-z>

Milani, D., Cross, J., Anderton, R., Blacker, D., Knuckey, N., & Meloni, B. (2017). Neuroprotective efficacy of poly-arginine R18 and NA-1 (TAT-NR2B9c) peptides following transient middle cerebral artery occlusion in the rat.

*Neuroscience Research*, 114, 9–15. <https://doi.org/10.1016/j.neures.2016.09.002>

Milani, D., Knuckey, N., Anderton, R., Cross, J., & Meloni, B. (2016). The R18 Polyarginine Peptide Is More Effective Than the TAT-NR2B9c (NA-1) Peptide When Administered 60 Minutes after Permanent Middle Cerebral Artery Occlusion in the Rat. *Stroke Research and Treatment*, 2016, 1–9. <https://doi.org/10.1155/2016/2372710>

Nijboer, C. H., van der Kooij, M. A., van Bel, F., Ohl, F., Heijnen, C. J., & Kavelaars, A. (2010). Inhibition of the JNK/AP-1 pathway reduces neuronal death and improves behavioral outcome after neonatal hypoxic-ischemic brain injury. *Brain, Behavior, and Immunity*, 24(5), 812–821. <https://doi.org/10.1016/j.bbi.2009.09.008>

Okusa, C., Oeschger, F., Ginet, V., Wang, W.-Z., Hoerder-Suabedissen, A., Matsuyama, T., ... Molnár, Z. (2014). Subplate in a rat model of preterm hypoxia-ischemia. *Annals of Clinical and Translational Neurology*, 1(9), 679–691. <https://doi.org/10.1002/acn3.97>

Ota, A., Ikeda, T., Ikenoue, T., & Toshimori, K. (1997). Sequence of neuronal responses assessed by immunohistochemistry in the newborn rat brain after hypoxia-ischemia. *American Journal of Obstetrics and Gynecology*, 177(3), 519–526. [https://doi.org/10.1016/S0002-9378\(97\)70139-X](https://doi.org/10.1016/S0002-9378(97)70139-X)

Palmer, C., Towfighi, J., Roberts, R. L., & Heitjan, D. F. (1993). Allopurinol administered after inducing hypoxia-ischemia reduces brain injury in 7-day-old rats. *Pediatric Research*, 33(4 Pt 1), 405–411. <https://doi.org/10.1203/00006450-199304000-00018>

Palmer, C., Vannucci, R. C., & Towfighi, J. (1990). Reduction of perinatal hypoxic-

ischemic brain damage with allopurinol. *Pediatric Research*, 27(4 Pt 1), 332–336.  
<https://doi.org/10.1203/00006450-199004000-00003>

Rao, R., Trivedi, S., Vesoulis, Z., Liao, S. M., Smyser, C. D., & Mathur, A. M. (2016). Safety and Short-Term Outcomes of Therapeutic Hypothermia in Preterm Neonates 34-35 Weeks Gestational Age with Hypoxic-Ischemic Encephalopathy. *The Journal of Pediatrics*. <https://doi.org/10.1016/j.jpeds.2016.11.019>

Rice, J. E., Vannucci, R. C., & Brierley, J. B. (1981). The influence of immaturity on hypoxic-ischemic brain damage in the rat. *Annals of Neurology*, 9(2), 131–141.  
<https://doi.org/10.1002/ana.410090206>

van den Broek, M. P. H., Groenendaal, F., Egberts, A. C. G., & Rademaker, C. M. A. (2010). Effects of Hypothermia on Pharmacokinetics and Pharmacodynamics. *Clinical Pharmacokinetics*, 49(5), 277–294. <https://doi.org/10.2165/11319360-000000000-00000>

Vaslin, A., Naegele-Tollardo, S., Puyal, J., & Clarke, P. G. H. (2011). Excitotoxicity-induced endocytosis mediates neuroprotection by TAT-peptide-linked JNK inhibitor. *J Neurochem*, 119(6), 1243–1252. <https://doi.org/10.1111/j.1471-4159.2011.07535.x>

Weinstock, M. T., Francis, J. N., Redman, J. S., & Kay, M. S. (2012). Protease-resistant peptide design-empowering nature's fragile warriors against HIV. *Biopolymers*, 98(5), 431–442. <https://doi.org/10.1002/bip.22073>

Wiegler, K., Bonny, C., Coquoz, D., & Hirt, L. (2008). The JNK Inhibitor XG-102 Protects from Ischemic Damage with Delayed Intravenous Administration Also in the Presence of Recombinant Tissue Plasminogen Activator. *Cerebrovascular Diseases*, 26(4), 360–366. <https://doi.org/10.1159/000151639>